



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07D 403/14, A61K 31/40, C07D 401/14, A61K 31/44</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/04732</b> <b>(43) International Publication Date:</b> 16 February 1995 (16.02.95)
<b>(21) International Application Number:</b> PCT/CA94/00427 <b>(22) International Filing Date:</b> 5 August 1994 (05.08.94) <b>(30) Priority Data:</b> 08/102,715                      6 August 1993 (06.08.93)                      US <b>(71) Applicant:</b> SYNPHAR LABORATORIES, INC. [CA/CA]; 4290 91A Street, Edmonton, Alberta T6V 5V2 (CA). <b>(72) Inventors:</b> LOWN, William, J.; 4704-117A Street, Edmonton, Alberta T6H 3SL (CA). MICETICH, Ronald, G.; 12 Braeside Terrace, Sherwood Park, Alberta T8A 3V6 (CA). <b>(74) Agent:</b> RICHES, McKENZIE & HERBERT; Suite 2900, 2 Bloor Street East, Toronto, Ontario M4W 3J5 (CA).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).  <b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> OLIGOPEPTIDE ANTIRETROVIRAL AGENTS		
$  \begin{array}{ccccccc}  \text{A}-(\text{NHCO})_x\text{-Het}-(\text{NHCO-Het})_y\text{-NH-R}_1\text{-NH}-(\text{Het-CONH})_z\text{-Het}(\text{CONH})_x\text{-A} & & & & & & \text{(I)} \\    & &   & &   & &   \\  \text{R}_2 & & \text{R}_3 & & \text{R}_4 & & \text{R}_5  \end{array}  $		
<b>(57) Abstract</b>  Oligopeptide antiretroviral agents are represented by formula (I) wherein A is a moiety bearing a positive charge and of a size which avoids steric inhibition of binding of said compound to nucleic acid sequences; R <sub>1</sub> is a moiety derived from a residue of carbonic acid or a residue of a dicarboxylic acid; Het is a five membered heterocyclic moiety; x is 0 or 1; y is 0, 1, 2 or 3; z is 0, 1, 2, or 3; R <sub>2</sub> , R <sub>3</sub> , R <sub>4</sub> and R <sub>5</sub> are attached to a ring atom other than carbon and are independently selected from the group consisting C <sub>1</sub> -C <sub>6</sub> alkyl and -CH <sub>2</sub> -O-R <sub>6</sub> , where R <sub>6</sub> is a C <sub>1</sub> -C <sub>6</sub> alkyl; or a salt thereof. These compounds exhibit antiretroviral activity, especially against HIV. These compounds are also useful against Hepatitis B.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## OLIGOPEPTIDE ANTIRETROVIRAL AGENTS


## FIELD OF THE INVENTION

This invention relates to oligopeptides which are particularly useful as antiretroviral agents.


## 5

## BACKGROUND OF THE INVENTION

Various oligopeptide derivatives have demonstrated various medicinal uses, such as enzyme inhibitors as disclosed in United States Patent 4,483,850. It is also known that various oligopeptides have anti-tumor activity as disclosed in United States Patents 4,216,208 and 4,314,999. Antibiotic activity of oligopeptides is disclosed in United States Patent 4,454,065. Naturally occurring oligopeptides, netropsin and distamycin, have been discovered as having antiviral and anti-tumor activity. The chemical formulas for netropsin and distamycin are as follows:



Compound 1



Compound 2

# SUBSTITUTE SHEET

These oligopeptides are disclosed in Julia, M., Préau-Joseph, N., C.R. Hebd-Seances, Acad. Sci. 1963, 257. 1115 and Arcamone, F.; Orezzi, P.G.; Barbier, W.; Nicolella, V.; Penco, S.; Gazz. Chim. Ital., 1967, 97, 1097.

Netropsin and distamycin contain pyrrole moieties connected by peptide bonds and with side chains, at least one of which is positively charged; i.e., an amidine group, or a group of the guanidyl type.

Only distamycin has been used as a therapeutic agent as commercialized and sold under the trade mark STALLIMYCIN HYDROCHLORIDE in the form of a 1% cream, ointment or paste. This composition has been used in the treatments of infections produced by herpes simplex, herpes zoster and vaccinia viruses. Topical application of distamycin has been limited due to its high cytotoxicity and a low therapeutic index which in the instance of treating the herpes virus is about 3.


U.S. Patent No. 4,912,199 discloses oligopeptides containing pyrrole moieties which demonstrated significantly enhanced antiviral and anticancer activities as compared to the oligopeptides of the prior art.

According to this invention oligopeptides have been developed which have significantly enhanced antiretroviral activity compared to prior types of oligopeptides.


#### SUMMARY OF THE INVENTION

According to an aspect of the invention, a compound represented by the formula I:

**SUBSTITUTE SHEET**




5



wherein A is a moiety bearing a positive charge and of a size which does not inhibit binding of said compound to nucleic acid sequences associated with the cellular action of retroviruses; R<sub>1</sub> is a moiety derived from a dicarboxylic acid or a residue of carbonic acid; Het is a five-membered heterocyclic moiety; y and z are independently 0, 1, 2 or 3, x is 0 or 1, and pharmaceutically acceptable salts thereof, exhibit antiretroviral activity, especially against Human Immunodeficiency Virus and Hepatitis B Virus.

A process for preparing such compounds comprises reacting a compound of the formula (II):



wherein x and y are as defined above; and B is the same as A or is a group with a nitrile, halogen or sulfide substituent; with a dicarboxylic acid of the formula (III):



wherein R<sub>1</sub> is as defined above and X is halogen, imidazolidine or other reactive moiety and converting B to A to form said moiety bearing a positive charge.

**SUBSTITUTE SHEET**

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph showing a correlation between DNA binding constants of linked oligopeptides ( $K_a$ ,-) and observed inhibitory properties expressed in reciprocal ID<sub>50</sub> values against Moloney Leukemia Virus reverse transcriptase (MlV-RT).

Figures 2-6 are graphs showing anti-HIV activity of several compounds of the present invention.



### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Compounds according to this invention demonstrate significant antiretroviral activity. Although the actual biological mechanism of these compounds which cause antiretroviral activity is not fully understood, it is thought that the activity may be due to the compounds of this invention binding with nucleic acid sequence(s) associated with the cellular action of retroviruses to inactivate such nucleic acids which code for the retroviral activity. It has also been observed that the linked oligopeptides of the present invention are potent inhibitors of Moloney Leukemia Virus (MIV) reverse transcriptase, a potential indicator of anti-HIV activity. See Figure 1.

The compounds of this invention have heterocyclic moieties, which may be the same or different, linked by a dicarboxylic acid derivative. Such linked heterocyclic moieties of this invention have significant unexpected activity compared to unlinked pyrrole moieties such as the naturally occurring netropsin and distamycin.

The compounds according to this invention are represented by the following formula:


**SUBSTITUTE SHEET**



wherein A is a moiety bearing a positive charge and of a size which does not inhibit binding of said compounds to deoxyribonucleic acid sequences associated with the cellular action of retroviruses;  $R_1$  is a moiety derived from a dicarboxylic acid; Het is a five-membered heterocyclic moiety;  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  may be attached to a ring carbon atom or hetero ring atom and are independently selected from  $C_1$ - $C_6$  alkyl and  $\text{CH}_2\text{-O-}R_6$ , where  $R_6$  is a  $C_1$ - $C_6$  alkyl;  $y$  and  $z$  are independently 0, 1, 2 or 3;  $x$  is 0 or 1; and pharmaceutically acceptable salts thereof.

The positively charged moiety at each extremity of the compound and identified as group A is preferably selected from the group of derivatives consisting of an amidine, a guanidine, secondary ammonium salts, sulfonium salts and phosphonium salts.


The selected amidine may have one or both nitrogen atoms of the amidine as a member of a five-membered cyclic structure. More particularly, the amidine derivative is represented by the formula:



wherein  $p$  equals 0 to 5 and  $X$  is  $-\text{H}$ ,  $-\text{OH}$ ,  $-\text{NH}_2$ ,  $-\text{CH}_3$ ,  $-\text{C}_2\text{H}_5$ ,  $-\text{C}_3\text{H}_7$ .


**SUBSTITUTE SHEET**

The selected guanidine for substituent A may be represented by the formula:




where p equals 0 to 5 and X equals -H, -OH, -NH<sub>2</sub>, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -C<sub>3</sub>H<sub>7</sub>.

10 When A is selected to be a quaternary, tertiary or secondary ammonium salt, it may be represented by the formula:



where p equals 1 to 5 and q equals 0 to 3 and X is an alkyl or alkenyl group of 1 to 3 carbon atoms.

15 When A is selected as a sulfonium salt, it may be represented by the formula:



where p equals 0 to 5 and X and Y are alkyl or alkenyl groups of 1 to 3 carbon atoms.

20 In the heterocyclic moieties, Het may be the same in each moiety or may be different. Preferably, the Het group is selected from the group consisting of a pyrrole, an imidazole, a triazole, a pyrazole, a thiazole, a thiophene, a furan, an oxazole and derivatives thereof.

25 Preferred ring carbon atoms substituents are alkyl groups, and especially methyl groups, on the Het moiety, especially on thiazole rings.


**SUBSTITUTE SHEET**



Preferred Het substituents are N-alkyl pyrrole having 1 to 6 carbon atoms in the alkyl group; N-alkyl imidazole having 1 to 6 carbon atoms in the alkyl group; alkyl pyrazole having 1 to 6 carbon atoms in the alkyl group; and alkyl triazole having 1 to 6 carbon atoms in the alkyl group. Preferably the N-alkyl pyrrole has 1 to 4 carbon atoms in the alkyl group, and especially in the N-methyl pyrrole. Also preferred Het substituents are N-linked alkoxymethyl groups. The choice of Het substituents will depend on their cellular uptake ability.

$R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are linked to the N or C atom of the Het moiety and are independently  $C_1$ - $C_6$  alkyl or  $-CH_2-O-R_6$  where  $R_6$  is  $C_1$ - $C_6$  alkyl. It has been found that the longer the alkyl group in either structure is, the better the cellular uptake of the compound. The choice of substituent will depend on solubility properties; solubility in pharmacologically acceptable solvents, such as water or DMSO, has been found to be higher with the methoxy substituents.

The linking group  $R_1$  is a derivative from carboxylic acid.  $R_1$  is represented generally by the formula:




where p equals any number from 1 to 22. Alternatively,  $R_1$

may be a residue of carbonic acid, namely  $-C-$ ; or  $R_1$  may be a residue of an aromatic dicarboxylic acid. The  $-CO-$  groups of the aromatic dicarboxylic acid residues may be in the ortho, meta or para positions on the ring. The aromatic residues may be 5 to 6 C membered rings. The aromatic dicarboxylic acid may also be a six membered heterocyclic ring containing a nitrogen atom.


## SUBSTITUTE SHEET

Other alternative structures for the linking group may be a residue of an unsaturated aliphatic dicarboxylic acid of the formula:



5 where q equals any number from 2 to 22.

$R_1$  may also be a residue of cycloalkane dicarboxylic acids of the formula:



10 where r equals any number from 3 to 7 and optionally may be fused to one or more three to seven C membered rings, preferably fused to one or two three to seven C membered rings.

$R_1$  may also be a residue of cycloalkane dicarboxylic acids of the formula:



where s equals any number from 3 to 7.

20 In a preferred compound of the present invention, A is a moiety selected from the group consisting of an amidine, a guanidine, secondary ammonium salts, tertiary ammonium salts, quaternary ammonium salts, sulfonium salts and phosphonium salts.


25 In another preferred compound of the present invention,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are each a  $\text{C}_1$ - $\text{C}_6$  alkyl or  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are the same and are a  $\text{C}_1$ - $\text{C}_6$  alkyl group or  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are each methoxymethyl.

**SUBSTITUTE SHEET**

In another preferred compound of the present invention,  $R_1$  is

5  $\begin{array}{c} \text{O} \\ || \\ -\text{C}- \end{array}$  or  $R_1$  is a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p\text{H}_{2p}-\text{CO}-$  where  $p$  equals 1 to 22.  $R_1$  may also be preferably a residue of a dicarboxylic acid selected from the group consisting of: a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where  $q$  equals 2; a residue of an aromatic dicarboxylic acid; and a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  where  $r$  equals 3 to 6.

In yet another preferred compound,  $R_1$  is



15 Preferably,  $R_1$  is a dicarboxylic acid residue of cyclopropane, a dicarboxylic acid residue of cyclopentane, or a dicarboxylic acid residue of cyclohexane.

The following are representative examples of the preferred compounds of the present invention.

20 N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] terephthalamide dihydrochloride.

N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] isophthalamide dihydrochloride.

25 N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] fumaramide dihydrochloride.


30 N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] maleamide dihydrochloride.

N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] trans 1,2-cyclobutanamide dihydrochloride.

**SUBSTITUTE SHEET**


N, N'-di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] trans 1,2-cyclobutanamide dihydrochloride.

The compound:




**SUBSTITUTE SHEET**

The compound:




**SUBSTITUTE SHEET**

The compound:




SUBSTITUTE SHEET

The compound:



SUBSTITUTED SHEET

The compound:



**SUBSTITUTE SHEET**







15

In cases where  $R_1$  is a dicarboxylic acid derivative of an aliphatic hydrocarbon, the linker is referred to as flexible. Rigid linkers refer to cases in which  $R_1$  is carbonic acid or residues of aromatic, unsaturated aliphatic, cycloalkane and cycloalkene dicarboxylic acids. Most preferred are those compounds in which  $R_1$  is a rigid linker. Examples of the flexible linked and rigid linked oligopeptides are set forth below.





















### FLEXIBLE LINKED OLIGOPEPTIDES

10





15


## SUBSTITUTE SHEET



# SUBSTITUTE SHEET


RIGID LINKED OLIGOPEPTIDES

5




19aR=R<sub>1</sub>  
19bR=R<sub>2</sub>


20aR=R<sub>1</sub>  
20bR=R<sub>2</sub>




trans 23aR=R<sub>1</sub>  
23bR=R<sub>2</sub>




cis 24aR=R<sub>1</sub>  
24bR=R<sub>2</sub>




trans 25aR=R<sub>1</sub>  
25bR=R<sub>2</sub>



trans 26aR=R<sub>1</sub>  
26bR=R<sub>2</sub>



trans 27aR=R<sub>1</sub>  
27bR=R<sub>2</sub>



cis 28aR=R<sub>1</sub>  
28bR=R<sub>2</sub>



10



21aR=R<sub>1</sub>  
21bR=R<sub>2</sub>


22aR=R<sub>1</sub>  
22bR=R<sub>2</sub>

**SUBSTITUTE SHEET**




SUBSTITUTE SHEET

Other preferred compounds include compounds of formula I wherein Het is pyrrole and x is 1; A is:




and  $R_1$  is a rigid linker (as defined above).

The heterocyclic moiety of the compounds of this invention may be linked in accordance with various processes by use of the dicarboxylic acid derivatives. In accordance with one aspect of this invention, the process for providing such linkage comprises reacting a compound of the formula:




wherein x and y are as defined above; and B is the same as A or is a group with a nitrile, halogen or sulfide substituent; with a dicarboxylic acid of the formula:



wherein  $R_1$  is as defined above and X is halogen, imidazolidine or other reactive moiety and converting B to A to form said moiety bearing a positive charge.

In the reactants, B may be generally represented by the formula:




where Z is CN-, hal or XS; hal is a halogen ion, X is an alkyl or alkenyl group having 1 to 3 carbon atoms, and p equals 0 to 5.

**SUBSTITUTE SHEET**

It is to be appreciated that B may also be identical to A in providing a charge group, for example, a guanidinium end group. In that instance, B has the general formula:

5




10

wherein X is an alkyl having 1 to 3 carbon atoms or alkenyl group having 2 or 3 carbon atoms and p equals 0 to 5 and s equals 0 or 1.


15

Compounds of the present invention which are asymmetrical around the linking group (i.e., wherein y and z are different in number) can be prepared by a two-step process, wherein the first step involves coupling a compound of the formula:




20

wherein B, x and z are as defined above, with a dicarboxylic acid of the formula:



25

wherein R<sub>1</sub> and X are as defined above (this coupling is generally with the use of equimolar amounts of the reactants). This is followed by coupling of a compound of the formula:




**SUBSTITUTE SHEET**


wherein B, x and z are as defined above, with the provisions that z is different than y.

According to preferred embodiments of the invention, the following reaction schemes demonstrate preferred chemical pathways to the compounds of this invention having the various desired end groups:


#### A - Preparation of Amidinium End Group




#### B - Preparation of Guanidinium End Groups



**SUBSTITUTE SHEET**

C - Preparation of Ammonium Salt in End Group


13

D - Preparation of Sulfonium Salts

SUBSTITUTE SHEET



Reference may be made to J.W. Lown and K. Krowicki, J. Org. Chem. 1985, 50, 3774 regarding the synthesis of related types of pyrrole moieties such as the synthesis of distamycin. The general synthesis of the compounds according to this invention are based on the total synthesis of distamycin. Dipyrrole or tripyrrole peptides bearing an amino group and a side-chain containing a group (B) which is the nitrile, ammonium or sulfide as represented by the following formula:



are allowed to react at a temperature of -35 to +10°C, preferably about -20°C, with a dicarboxylic acid dichloride in the presence of a base or with a diimidazolidine of a dicarboxylic acid to give a bis-amide of the dicarboxylic acid. The resulting compound in the case of nitrile is allowed to react at a temperature of 0 to +35°C, preferable +15° to +25°C, more preferably about +20°C, with ethanol in the presence of hydrochloric acid and then at a temperature of 0 to +35°C, preferably +15 to +25°C, more preferably about +20°C, with ammonia (Pinner reaction) to generate an amidinium moiety in the final product, as exemplified by the above reaction scheme A. As with reaction scheme D, the sulfide is methylated at a temperature of 0 to +35°C, preferable +15 to +25°C, more preferably about +20°C, to produce the corresponding sulfonium salt.

The compounds of formula I, are useful as antiretroviral agents, especially against the Human Immunodeficiency Virus (HIV). Human patients suffering

**SUBSTITUTE SHEET**

from diseases caused by , for example, HIV, can be treated by administering to the patient a pharmaceutically effective amount of one or more of the present compounds optionally, but preferably in the presence of a  
5 pharmaceutically acceptable carrier or diluent. There may be also included pharmaceutically compatible binding agents and/or adjuvant materials. The active materials can also be mixed with other active materials which do not impair the desired action and/or supplement the desired  
10 action. The active materials according to the present invention can be administered by any route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, rectally or topically, in a liquid or solid form. For injection purposes, the medium used may  
15 be a sterile liquid. As an injection medium, it is preferred to use water which contains the stabilizing agents, solubilizing agents and/or buffers conventional in the case of injection solutions. Desirable additives include, for example, tartrate and borate buffers,  
20 ethanol, dimethylsulfoxide, complex forming agents (for example, ethylenediamine tetracetic acid), high molecular weight polymers (for example, liquid polyethylene oxide) for viscosity regulation or polyethylene derivatives or sorbitan anhydrides. Solid carrier materials include, for  
25 example, starch, lactose, mannitol, methylcellulose, talc, highly dispersed silicic acid, high molecular weight fatty acids (such as stearic acid), gelatin, agar, calcium phosphate, magnesium stearate, animal and vegetable fats of solid high molecular weight polymers (such as polyethylene glycol).  
30 Compositions suitable for oral administration can, if desired, contain flavoring and/or sweetening agents.

A preferred mode of administration of the compounds of this invention is oral. Accordingly, the  
35 compounds may be formulated into capsule form or tablet form.

**SUBSTITUTE SHEET**

The active materials according to the present invention can be employed in dosages and amounts which are conventional in the art. Thus, the materials can be used at a dosage range in humans of from about 1 to 200 mg/kg total body weight/day. A more preferred range lies between 1-30 mg/kg total body weight/day. The dosages may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

The in vitro anti-HIV screening test results, performed at the United States National Cancer Institute, have shown that 23 of the present compounds are active. Of the fifteen, ten are considered "active", and thirteen are determined "moderately active". Certain of the compounds screened for anti-AIDS activity at the NCI were determined to be "inactive". These compounds were ones wherein the  $R_1$  is  $-\text{CO}-(\text{CH}_2)_6-\text{CO}-$  or  $-\text{CO}-(\text{CH}_2)_8-\text{CO}-$ , A is amidine, x is 1, Het is methylpyrrole, and y and z are 1, as well as compounds 9, 11, 15, 16, 18 and 37.

The therapeutic index of a compound is determined by dividing the inhibitory or lethal concentration for 50% of the population ( $\text{IC}_{50}$ ) by the effective concentration for 50% of the population ( $\text{EC}_{50}$ ). The therapeutic indexes for the particularly active compounds of the present invention range from 1.46 to 161.

As used in this invention, antiretroviral activity refers to the ability of a compound to inhibit the growth of a retrovirus. The retrovirus of primary importance with respect to the present invention is HIV. However, the present compounds may also exhibit antiretroviral activity towards other retroviruses as would be apparent by the suspected mechanism of action and other viruses which replicate or exhibit reverse transcription.

**SUBSTITUTE SHEET**

The compounds of the present invention should also be therapeutically effective in the treatment of hepatitis B viral infection in mammals, especially humans. Similar to retroviruses (including HIV-1), the hepatitis B virus replicates by reverse transcription. In addition, hepatitis B virus putative viral polymerase share amino acid homology with reverse transcriptase of retroviruses and a comparison of the thirteen (13) hepadnavirus isolates determined that other conserved areas showing homology to corresponding regions of Type C retro virus. Miller et al., Proc. Natl. Acad. Sci. USA, Vol 83:2531-2535 (1986).

Since it is theorized that the activity of the compounds of the present invention may be due to the compounds binding with nucleic acid sequence(s) associated with the cellular action of retroviruses to inactivate such nucleic acids which code for the retroviral activity, the compounds are likely to inhibit binding with nucleic acid sequence(s) of the hepatitis B virus associated with the cellular action of reverse transcription to inactivate such nucleic acids which code for the retroviral-like activity. Therapeutically effective anti-hepatitis B dosages would be the same as anti-HIV-1 dosage levels as well as would the routes of administration.


The ability of a compound to inhibit HIV may be measured by various experimental techniques. One such technique, currently employed by the United States National Cancer Institute to screen potential anti-HIV compounds, involves the inhibition of the killing of HIV-infected T<sub>4</sub> lymphocytes. Compounds of the present invention have been tested for anti-HIV-1 activity in the NCI protocol; however, one skilled in the art would appreciate that the compounds should exhibit activity against HIV-2 as well.

**SUBSTITUTE SHEET**

Preferred embodiments of the invention are exemplified in the following Examples which are in no way to be construed as limiting the scope of the appended claims.

5 EXAMPLE 1

Compounds of the formula I, where x=1, y and z each are 1;




$R_1$  equals  $-\text{COCH}_2\text{CH}_2\text{CO}-$ , was prepared. 1-Methyl-4-(1-methyl-4-aminopyrrole-2-carboxamido)-pyrrole-2-carboxamidopropionitrile (105 mg, 0.33 mmole) and 1-Pr<sub>2</sub>EtN (diisopropylethylamine) (65  $\mu$ l, 0.16 mmole) in anhydrous THF (1 ml) was added and the mixture was allowed to reach room temperature. The solvents were evaporated to dryness and water was added. The resulting solid was collected and washed with hot MeOH to give 90 mg (77% yield) of the product m.p. 297°C. The latter was suspended in anhydrous EtOH and saturated with HCl while cooling. After 1.5 hours at room temperature, the solvent was removed in vacuo and the residue was washed with dry ether then ethanol was added followed by some ammonia condensed into the solution. After 1 hour at room temperature, the solvent was removed and the residue was washed with MeOH, EtOH and hexane to afford 80 mg of a solid. Recrystallization from a small volume of water gave a jelly-like precipitate which was washed with EtOH, hexane and dried to give 35 mg (35% yield) of pure product m.p. 283-285°C dec. <sup>1</sup>H-NMR (DSMO-d<sub>6</sub>):  $\delta$  2.60 (m, 4H), 3.60 (m,

# SUBSTITUTE SHEET

2H), 3.83 (s, 6H), 6.92 (d, 2H), 7.18 (d, 2H), 8.25 (t, 1H), 8.70 (bs, 2H), 9.02 (bs, 2H), 9.93 and 9.97 (2s, 2H), MS-FAB (m/z): 745 (M-Cl-HCl)<sup>+</sup>; Anal. Calcd. for C<sub>34</sub>H<sub>46</sub>Cl<sub>2</sub>H<sub>14</sub>O<sub>6</sub>: C, 49.9, H, 5.7, N, 24.0, Cl, 8.7, Found: 5 C, 50.3, H, 6.05, N, 22.9, Cl, 8.7.

EXAMPLE 2

Compounds of the formula I, where x equals 1; y and z each are each equal to 1;



R<sub>1</sub> equals -CO- was prepared. 1-Methyl-4-(1-methyl-4-aminopyrrole-2-carboxamido)-pyrrole-2-carboxamidopropionitrile (315 mg, 1 mmole) and 81 mg of 15 1,1'-carbonyldiimidazole were dissolved in 10 ml of anhydrous CH<sub>3</sub>CN and refluxed under argon for 5 minutes. A solid forms which was collected to give 302 mg (88.6% yield) of the pure product was treated with HCl in EtOH and then NH<sub>3</sub> (as in Example 1). After the reaction was 20 completed, the mixture was decanted from an insoluble residue. The solvent was removed in vacuo and the residue was dissolved in 4 ml of MeOH and an excess of CH<sub>3</sub>CN was added to precipitate the product which was collected and washed with 1 ml of cold water whereupon it became jelly-like. 25 The product was redissolved in MeOH and reprecipitated with CH<sub>3</sub>CN to give 216 mg (57% overall yield) of the pure compound m.p. 211-215°C; <sup>1</sup>H-NMR (DMS-d<sub>6</sub>): δ 2.64 (t, 2H), 3.52 (q, 2H), 3.84 (s, 6H), 6.82, 6.94, 7.03, 7.20 (4d, 4H), 8.25 (t, 1H), 8.73 (2s, 3H), 30 9.05 (s, 2H), 9.88 (s, 1H), MS-FAB: 690 (M-Cl-HCl)<sup>+</sup>. Anal. Calcd. for C<sub>31</sub>H<sub>42</sub>Cl<sub>2</sub>N<sub>14</sub>O<sub>5</sub>: C, 48.9, H, 5.6, Cl, 9.3, N, 25.7; Found: C, 48.5, H, 5.7, Cl, 9.7, N, 25.3.

**SUBSTITUTE SHEET**

EXAMPLES 3(A) AND (B)

5 The following Examples illustrate the effect of altering the steric size of the terminal group (in these cases trialkylammonium) on the basic Het block of the general formula on the nucleic acid binding and antiviral efficacy. The effects were demonstrated on deoxyribonucleic acid to show that steric hindrance in the terminal group in DNA binding generally reduces antiviral activity of the compounds.

- 10 (A) 1-Methyl-4-(1-methyl-4-trimethylammonium-acetamidopyrrole-2-carboxamido)pyrrole-2-carboxyamidopriopionamidine chloride hydrochloride

15 A solution of the precursor 1-methyl-4-(1-methyl-4-trimethylammonium-acetamido-pyrrole-2-carboxamido)pyrrole-2-carboxyamidopriopionitrilechloride (347 mg, 0.07 mmoles) in 5 ml of absolute ethanol was treated with dry hydrogen chloride with cooling. After 2 hours, the solvent was removed in vacuo, 5 ml of absolute ethanol was added and dry NH<sub>3</sub> gas passed into the solution. The solid dissolved during 2 hours at room temperature, then the solution was evaporated to dryness and extracted with hot isopropyl alcohol (100 ml). The extract was concentrated to ca. 10 ml, acetone added and the resulting precipitate collected, washed with acetone, and dried to vacuo to give the product, 300 mg (85% yield) as an amorphous hygroscopic solid, no definite m.p.; <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.67 (t, 2H), 3.31 (s, 9H), 3.52 (q, 2H), 3.82 and 3.87 (2s, 6H), 4.44 (s, 2H), 6.97 (d, 1H), 7.02 (d, 1H), 7.24 (d, 1H), 7.29 (d, 1H), 7.29 (d, 1H), 8.31 (t, 1H), 8.82 (bs, 2H), 9.72 (bs, 2H), 10.06 (s, 1H), 11.23 (s, 1H), IR (Nujol) ν<sub>max</sub>: 1260, 1377, 1405, 1453, 1531, 1582, 1643, 1685, 3247 cm<sup>-1</sup>; MS-FAB (m/z) 430 (M-HCl-Cl)<sup>+</sup>.

**SUBSTITUTE SHEET**

Sulfate. The sulfate corresponding to the product was prepared in order to obtain an analytically pure sample by precipitation from a methanolic solution of the above compound by means of a large excess of tetraethylammonium sulfate, m.p. 295°C: IR (Nujol)  $\nu_{\max}$ : 1255, 1377, 1405, 1462, 1525, 1560, 1580, 1640, 1670, 3280  $\text{cm}^{-1}$ ; MS-FAB (m/z) 431 ( $\text{M-HSO}_4$ )<sup>+</sup>, 529  $\text{MH}^+$ ; Anal. Calcd. for  $\text{C}_{20}\text{H}_{32}\text{N}_8\text{O}_7\text{S}$  (528.59), C, 45.4, H, 6.1, N, 21.1, S, 6.1. Found: C, 45.0, H, 6.0, N, 20.7, S, 5.8.

- 10 (B) 1-Methyl-4-(1-methyl-4-trimethylammonium-acetamidopyrrole-2-carboxamido)pyrrole-2-carboxyamidopriopionamidine chloride hydrochloride

A solution of the precursor analogous to that of Example 3(A) (173 mg, 035 mmoles) in 10 ml of absolute ethanol was treated with dry hydrogen chloride with cooling. After 2 hours, the solvent was removed in vacuo and the residue dissolved in 10 ml of absolute ethanol and treated with an excess of dry ammonia. After 2 hours at room temperature, the solvent was removed in vacuo and the residue dissolved in 5 ml of isopropyl alcohol; then the product was precipitated with ether. The solid was collected, washed with ether and dried at 100° in vacuo to afford the product 103 mg (59% yield) m.p. 180° (dec); <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>):  $\delta$  1.32 (t, 9H), 2.67 (t, 2H), 3.54 (m, 8H), 3.83 and 3.88 (2s, 6H), 4.32 (s, 2H), 6.96 (d, 1H), 7.01 (d, 1H), 7.21 (d, 1H), 7.30 (d, 1H), 8.28 (t, 1H), 7.01 (d, 1H), 7.21 (d, 1H), 7.30 (d, 1H), 8.28 (t, 1H) 8.80 and 9.10 (bs, 4H), 10.03 (s, 1H), 11.47 (s, 1H), IR (Nujol): 1367, 1404, 1462, 1531, 1581, 1646, 1684, 3250  $\text{cm}^{-1}$ ; MS-FAB (m/z): 981 (2M-HCl-Cl)<sup>+</sup>, 473 (M-HCl-Cl)<sup>+</sup>.

**SUBSTITUTE SHEET**



The activities of Examples 3(A) and 3(B) expressed as minimum inhibitory concentration ( $\mu\text{g/ml}$ ) against vaccinia virus were 20 and 300 respectively illustrating the effects of steric hindrance in DNA binding on reducing agent activity. The larger the terminal group, as demonstrated by compound 3(B), the lesser the activity; hence the terminal group is of a selected size which will maintain nucleic acid sequence bonding desired antiretroviral activity.

The compound numbers referred to in the following examples correspond to the numbered structures in the "Detailed Description of the Invention" section.

#### EXAMPLE 4

(A) 1-Methyl-4-[1-methyl-4-(1-methyl-4-aminopyrrole-2-carboxamido)pyrrole-2-carboxamido]pyrrole-2-carboxamidopriopionitrile (Intermediate Compound)

1-Methyl-4-[1-methyl-4-(1-methyl-4-aminopyrrole-2-carboxamido)pyrrole-2-carboxamido]pyrrole-2-carboxamidopriopionitrile (Lown, W.J. and Krowicki, K., J. Org. Chem. Vol. 50, p. 3774 (1985) and Krowicki, K. and Lown, W.J., J. Org. Chem., Vol. 52, p. 3493 (1987) (420 mg, 0.9 mmol) was reduced over 5% palladium on charcoal (260 mg) in a mixture of DMF (15 ml) and methanol (5 ml) at 45°. After the reduction the solvents were evaporated under reduced pressure. The residue was dissolved in a small amount of acetonitrile (2 ml) and an excess of ethyl acetate (20-30 ml) as added to precipitate some impurities. The filtrate was treated with an excess of hexane to precipitate a white pure product 9 (250 mg, 63.5% yield), m.p. 155-160°.  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ):  $\delta$  2.74 (t, 2H), 3.42 (q, 2H), 3.76 (s, 3H), 3.85 and 3.87 overlapped with a bs (3s, 8H), 6.27 (d, 1H), 6.40 (d, 1H), 6.95 (d, 1H), 7.04 (d, 1H), 7.24 (2d, 2H), 8.37 (t, 1H),

**SUBSTITUTE SHEET**

9.66 (s, 1H), 9.96 (s, 1H); IR (nujol): 1260, 1377, 1403, 1464, 1529, 1582, 1646, 2245, 3120, 3310  $\text{cm}^{-1}$ ; MS m/z 436.1981 (calcd. 436.1983). Analysis Calcd. for  $\text{C}_{19}\text{H}_{26}\text{ClN}_6\text{O}_3$ : C 52.3, H 6.0, Cl 8.1, N 22.5. Found: C 52.3, H 6.0, Cl 7.9, N 22.0.

#### EXAMPLE 5

(A) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido(3-proprionitrile)-4-pyrrole]-4-pyrrolyl)succinamide

The intermediate compound (105 mg, 0.33 mmol) and  $i\text{-Pr}_2\text{EtN}$  (65  $\mu\text{L}$ , 0.37 mmol) were dissolved in anhydrous acetonitrile (5 ml) and cooled to  $-20^\circ\text{C}$ . Succinyl chloride (18  $\mu\text{L}$ , 0.16 mmol) in anhydrous THF (1 ml) was added. The mixture was allowed to reach ambient temperature. The solvents were evaporated to dryness, water was added, and the resulting solid was collected and washed with hot MeOH. The product was dissolved in DMF and when placed on a TLC plat ( $\text{SiO}_2$ ) with  $\text{CHCl}_3$  + 15% MeOH system it gave one spot. For analytical purposes, the product was purified by dissolution in a small amount of DMF and precipitation was large amount of EtOH to give 90 mg (77%) of 15 m.p.  $292^\circ$ .  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ):  $\delta$  2.58 (s, 4H) 2.74 (t, 4H), 3.42 (q, 4H), 3.83 (2s, 12H), 6.86, 6.93, 7.17 and 7.22 (4d, 2H each), 8.35 (t, 2H), 9.89 (s, 4H); IR (nujol): 1376, 1401, 1447, 1465, 1511, 1535, 1585, 1645, 2245, 3120, 3304  $\text{cm}^{-1}$ ; MS (m.z. rel. int.): 396.1543 (9.98) for  $\text{C}_{19}\text{H}_{20}\text{N}_6\text{O}_4$  which is  $(\text{O}=\text{C}=\text{CH}-\text{M}_{1/2})^+$ . Analysis Calcd. for  $\text{C}_{34}\text{H}_{38}\text{N}_{12}\text{O}_6$ : C 57.5, H 5.4, N 23.6. Found: C 57.8, N 5.4, N 23.3.

**SUBSTITUTE SHEET**

(B) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido  
(3-proprionamidine)-4-pyrrole]-4-pyrrolyl)succinamide  
dihydrochloride (Compound 5)

5 A suspension of the previous product (130 mg,  
0.18 mmol) in 15 ml anhydrous EtOH was saturated with HCl  
with cooling. After 1.5 hr. at r.t., the solvent was  
evaporated under reduced pressure. The residue was washed  
with dry ether, then ethanol was added followed by some  
NH<sub>3</sub> condensed into the vessel. After 1 hr at r.t. the  
10 solvents were removed and the residue was washed with  
MeOH, EtOH and hexane to give 116 mg of a solid. The  
latter was examined by TLC (SiO<sub>2</sub>) with MeOH and a drop of  
formic acid and indicated formation of the product  
(R<sub>f</sub> = 0.3) containing some more polar impurity.  
15 Recrystallization from a small amount of water gave a gel-  
like precipitate which was washed with EtOH and hexane and  
dried give to 50 mg (34% of pure 5a, m.p. 283-5° dec.  
<sup>1</sup>H=NMR (DMSO-d<sub>6</sub>): δ 2.60 (m, 8H) 3.50 (m, 4H), 3.83  
(s, 12H), 6.92 (d, 4H), 7.18 (d, 4H), 8.25 (t, 2H), 8.70  
20 (bs, 4H), 9.02 (bs, 4H) 9.93 and 9.97 (2s, 4H); IR  
(nujol): 1352, 1377, 1464, 1521, 1576, 1638, 1700, 3260  
cm<sup>-1</sup>; MS-FAB (m/z): 745 (M-Cl-HCl)<sup>+</sup>. Analysis Calcd. for  
C<sub>34</sub>H<sub>46</sub>Cl<sub>2</sub>N<sub>14</sub>O<sub>6</sub>: C 49.94, H 5.67, N 23.98, Cl 8.67. Found:  
C 50.3, H 6.05, N 22.90, Cl 8.75.

25 EXAMPLE 6

(A) N, N'-Di(1-methyl-2-[1-methyl-2-carboxamido  
(3-proprionitrile)-4-pyrrole]-4-pyrrolyl)malonamide

The intermediate compound (315 mg, 1 mmol),  
malonic acid (52 mg, 0.5 mmol) and DCC 206 mg, 1 mmol)  
30 were stirred in acetonitrile (6 ml) for 2 hr at room  
temperature and finally the mixture was heated briefly to

**SUBSTITUTE SHEET**

boiling to complete the reaction. A solid which contained dicyclohexylurea was collected and the filtrate was extracted with DMF. The DMF solution was treated with water and the solid formed was recrystallized from a mixture of acetonitrile (2 ml) and methanol (2 ml) to give pure compound (140 mg, 40% yield), m.p. 225-30°. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.73 (t, 2x2H), ~2.40 (q+s overlapped, 2x2H+2H), 3.83 and 3.86 (2s, 2x6H), 6.91 (2d, 2x2H), 7.18 and 7.22 (2d, 2x2H), 8.35 (t, 2x1H), 9.91 (s, 2x1H), 10.09 (s, 2x1H); IR (nujol): 1200, 1264, 1290, 1376, 1401, 1464, 1511, 1532, 1585, 1638, 1662, 2250, 3120, 3305 cm<sup>-1</sup>; MS-FAB (m/z): 697 (MH<sup>+</sup>). Analysis Calcd. for C<sub>33</sub>H<sub>36</sub>N<sub>12</sub>O<sub>6</sub>: C 56.9, H 5.2, N 24.1. Found: C 56.6, H 5.4, N 23.9.

(B) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido (3-proprionamide)-4-pyrrole]-4-pyrrolyl}malonamide dihydrochloride (Compound 4)

The compound of the previous synthesis (160 mg, 0.23 mmol) was suspended in dry ethanol and the mixture was saturated with dry halogen chloride. After 1.5 hr at room temperature, the solvent was removed under reduced pressure. The residue was treated with dry ethanol and dry ammonia. After 1 hr the solution was decanted from undissolved material and evaporated to dryness. The residue was dissolved in 2 ml of boiling water and an excess of acetonitrile was added to the hot solution. The precipitate was collected and washed with a small amount of water. The operation was repeated and pure compound 4 was collected, 100 mg (59% yield), m.p. 218-224°. The compound, if crystallized from water, precipitates in the form of a jelly. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.63 (t, 2x2H), ~3.35 (s overlapped with the peak of water), 3.50 (q, 2x2H), 3.80 and 3.83 (2s, 2x6H), 6.93 (s, 2x2H), 7.20

**SUBSTITUTE SHEET**

(s, 2x2H), 8.26 (t, 2x1H), 8.90 (bs, 2x4H), 9.96 (s, 2x1H), 10.28 (s, 2x1H). D<sub>2</sub>O exchange experiment showed the presence of malonyl protons at  $\delta$ 3.30. IR (nujol): 1260, 1377, 1405, 1463, 1535, 1580, 1645, 3100, 3270 cm<sup>-1</sup>; MS-FAB (m/z) 731 (M-Cl-HCl)<sup>+</sup>. Analysis Calcd. for C<sub>33</sub>H<sub>44</sub>N<sub>14</sub>O<sub>6</sub>Cl<sub>2</sub>: C 49.3, H 5.5, N 24.4, Cl 8.8. Found: C 49.0, H 5.7, N 27.0, Cl 9.0.

#### EXAMPLE 7

(A) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido  
10 (3-proprionitrile)-4-pyrrole]-4-pyrrolyl}urea

The intermediate compound (365 mg, 1.16 mmol) and 1,1'-carbonyldiimidazole (94 mg, 0.58 mmol) were allowed to react in boiling acetonitrile (3 ml). A solid which formed was collected, washed with acetonitrile to  
15 give 350 mg (88.6% yield) of pure product, m.p. 296-7°. <sup>1</sup>H-NMR 3.88 (s, 6H), 6.80 (d, 2H), 6.92 (d, 2H), 7.02 (d, 2H), 7.21 (d, 2H), 8.12 (s, 2H), 8.25 (t, 2H), 9.81 (s, 2H); IR (nujol): 1199, 1217, 1252, 1378, 1409, 1436, 1465, 1504, 1544, 1589, 1621, 1653, 1672, 2240, 3270, 3424 cm<sup>-1</sup>;  
20 MS-FAB (m/z): 655 (MH<sup>+</sup>). Analysis Calcd.: C 56.9, H 5.2, N 25.7. Found: C 56.6, H 5.4, N 25.5.

(B) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido  
(3-proprionamide)-4-pyrrole]-4-pyrrolyl}urea  
dihydrochloride (Compound 3)

25 The compound synthesized in the previous step (116 mg, 0.25 mmol) was suspended in dry ethanol and the solution saturated with HCl. After 2 hr the solvent was evaporated in vacuo and the residue treated with dry ammonia in ethanol for 1 hour. The mixture was decanted  
30 from an insoluble residue and the solution evaporated to

**SUBSTITUTE SHEET**

dryness. The residue was dissolved in 2 ml of methanol and an excess of acetonitrile was added to precipitate the product. The latter was collected and washed with 1 ml of water when it became jelly-like. It was redissolved in methanol and precipitated with acetonitrile to give the compound (3) (117 mg, 61.6% yield), m.p. 211-215°. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.64 (t, 4H), 3.52 (q, 4H), 3.84 (2s, 12H), 6.82 (d, 2H), 6.94 (d, 2H), 7.03 (d, 2H), 7.20 (d, 2H), 8.73 (2s overlapped, 6H), 9.05 (s, 4H), 9.88 (s, 2H); IR (Nujol): 1264, 1377, 1402, 1439, 1489, 1531, 1583, 1640, 1689, 3088, 3279 cm<sup>-1</sup>; MS-FAB (m/z): 690 (M-Cl-HCl)<sup>+</sup>. Analysis Calcd. for C<sub>31</sub>H<sub>42</sub>Cl<sub>2</sub>N<sub>14</sub>O<sub>5</sub>: C 48.9, H 5.6, Cl 9.3, N 25.7. Found: C 48.5, H 5.7, Cl 9.7, N 25.3.

#### EXAMPLE 8

(A) N,N'-Di{1-methyl-2-carboxamido(3-proprionitrile)-4-pyrrolel-4-pyrrolyl}adipamide

Adipic acid (29.2 mg, 0.2 mmol) in acetonitrile (0.5 ml) was treated with pivaloyl chloride (50 μL, 0.4 mmol) and Hunig's base (160 μL, 0.9 mmol) and then compound 7 (126 mg, 0.42 mmol) in DMF (0.5 ml) was added. After a half hour at room temperature the mixture was evaporated to dryness under reduced pressure. The residue was washed with water and hot acetonitrile. The solid was dissolved in hot DMF and precipitated with an excess of acetonitrile to give the compound (95 mg, 61% yield), m.p. 244-46° dec. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.60 (s, 4H), 2.27 (s, 4H), 2.74 (t, 4H), 3.40 (q, 4H), 3.83 (2s, 12H), 6.86 (s, 2H), 6.93 (s, 2H), 7.17 (s, 2H), 7.22 (s, 2H), 8.38 (t, 2H), 9.82 (s, 2H), 9.91 (s, 2H); IR (Nujol): 1376, 1400, 1464, 1513, 1533, 1585, 1641, 2258, 3294 cm<sup>-1</sup>; MS-FAB (m/z): 738 (M<sup>+</sup>), 739 (MH<sup>+</sup>); Analysis Calcd. C 58.5, H 5.7, N 22.7. Found: C 58.9, H 5.9, N 22.5.

**SUBSTITUTE SHEET**

(B) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-pyrrolyl}adipamide dihydrochloride (Compound 7)

The compound synthesized in the previous step  
5 (320 mg, 0.43 mmol) was treated under Pinner reaction conditions as in Example 3 above. After evaporation of solvents, water (3.5 ml) was added and a crystalline substance was collected to give (7) (215 mg, 58.7% yield),  
10 m.p. 195-6°. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.60 (s, 4H), 2.27 (s, 4H), 2.62 (t, 4H), 3.52 (q, 4H), 3.80 (2s, 12H), 6.88 (d, 2H), 6.95 (d, 2H), 7.18 and 7.20 (2d, 4H), 8.25 (t, 2H) 8.70 (s, 4H), 9.00 (s, 4H), 9.00 (s, 4H), 9.92 (s, 2H); IR (Nujol): 1208, 1261, 1377, 1404, 1463, 1531, 1579, 1641, 1691, 3256 cm<sup>-1</sup>; MS-FAB m/z 773 (M-HCl-Cl)<sup>+</sup>;  
15 Analysis Calcd.: C 51.1, H 6.0, N 23.2, Cl 8.4. Found: C 50.9, H 6.2, N 23.6, Cl 8.8.

#### EXAMPLE 9

(A) N,N'-Di(1-methyl-2-[1-methyl-2-carboxamido(3-propionitrile)-4-pyrrole]-4-pyrrolyl}maleimide

20 The intermediate compound (158 mg, 0.5 mmol) and maleic anhydride (49 mg, 0.5 mmol) were heated in acetonitrile (5 ml) at 50° for 3 minutes. Another portion of the intermediate compound (158 mg) was added and the solution was evaporated to dryness. The residual solid  
25 was dissolved in DMF (2 ml) and DCC (103 mg, 0.5 mmol) was added, and the mixture was set aside overnight at room temperature. Two drops of water were added and the solution was filtered. Then an excess of water precipitated the crude product. The product was collected  
30 and chromatographed on silica gel with chloroform and 15% of methanol providing yellow fractions. These were combined and evaporated, and the residue recrystallized

**SUBSTITUTE SHEET**

from acetone to give the product (100 mg, 56.5% yield), m.p. 250-2°. Analytical data for this and related compounds is given in Table I.

5 (B) N,N'-Di{1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole}-4-pyrrolyl}maleamide dihydrochloride  
Compound (14)

10 The compound obtained in the previous step (170 mg, 0.24 mm) was treated under Pinner reaction conditions as in Example 3. The completed reaction mixture was  
15 evaporated to dryness and the residue dissolved in ethanol. Controlled addition of isopropanol provided selective precipitation of impurities. The mother liquor was evaporated and the residue was dissolved in methanol and precipitation with acetonitrile gave pure compound  
(14) (166 mg, 85% yield), m.p. 217°.

#### EXAMPLE 10

(A) N,N'-Di{1-methyl-2-[1-methyl-2-carboxamido(3-propionitrile)-4-pyrrole]-4-pyrrolyl}trans  
cyclopropyldicarboxamide (Compound 8a)

20 The synthesis and characterization of compounds 3, 4 and 5 have been reported (Krowicki, K. et al, J. Med. Chem., Vol. 31, p. 341 (1988)). Trans-cyclopropyl-  
dicarboxylic acid (59 mg, 0.45 mmole) and 1,1'-  
25 carbonyldiimidazole (146 mg, 0.7 mmole) in acetonitrile (2.5 ml) were heated under reflux until the evolution of carbon dioxide ceased. To the cooled solution the  
appropriate amine (284 mg, 0.9 mmole) and 0.8 ml of DMF  
30 were added and the mixture was stirred for 2 hr at room temperature (the product partially precipitated) and was evaporated to dryness under reduced pressure. The residue was washed with acetonitrile, aqueous K<sub>2</sub>CO<sub>3</sub> then water to give 8a, 289 mg (88.6% yield) m.p. 312° dec.

**SUBSTITUTE SHEET**



(B) N,N'-Di{1-methyl-2-[1-methyl-2-carboxamide(3-proprionamidine)-4-pyrrole]-4-pyrrolyl} trans-cyclopropyldicarboxamide dihydrochloride  
(Compound 8b)

---

5                   Compound 8a (216 mg, 0.3 mmole) was treated  
under Pinner reaction conditions as described previously.  
The final reaction mixture was evaporated to dryness and  
the residue was extracted with hot propanol (150 ml). The  
extract was evaporated to dryness and the residue  
10 dissolved in methanol 1 ml, and an excess of acetonitrile  
was added to precipitate the product 8b, 170 mg  
(68.5% yield) m.p. 210° (softens).

#### EXAMPLE 11

15                   Commercially available acid chlorides for the  
linker groups were used directly without further  
purification. Otherwise, the appropriate acid chlorides  
were prepared from the acids according to the following  
procedure: An acid and a drop of dimethylformamide was  
heated in thionyl chloride (5 to 10 mole in excess) to 55-  
20 65°C for 30 to 45 min until a homogeneous liquid was  
obtained. The excess of the chlorinating agent was  
removed by evaporation. A small amount of methylene  
chloride was added to the crude acid chloride then  
evaporated. The diacid dichloride was then dissolved in  
25 methylene chloride or THF and aliquots were taken and used  
for coupling reactions.

#### EXAMPLE 12

30                   Distamycin A (50 mg, 0.09 mmol) was dissolved in  
4 mL of methanol. To this yellow solution was added 100  
μL of concentrated hydrochloric acid. The solution was  
stirred for 6-8 h and the reaction progress was followed  
by TLC (methanol:acetic acid, 100:5). The solvent was

**SUBSTITUTE SHEET**

evaporated and the crude product was redissolved in methanol and precipitated with ether. The product was recrystallized in this way twice more. The supernatant was decanted and the residual solid was dried in vacuo. The final product was obtained as an off-white solid 50 mg (89% yield).

### EXAMPLE 13

#### Bis-distamycin (Compound 15)

A solution of succinyl dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 68% yield. m.p. 210°C;  $^1\text{H-NMR}$ , 2.48 (COCH<sub>2</sub>CH<sub>2</sub>CO, 4H, s), 2.56 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6Hz], 3.50 (2xCONHCH<sub>2</sub>, 4H, q, J = 6Hz), 3.80 (2xNCH<sub>3</sub>, 6H, s), 3.82 (2xNCH<sub>3</sub>, 6H, s), 3.83 (2xNCH<sub>3</sub>, 6H, s), 6.90 (2xpy-CH, 2H, d, J = 2 Hz), 6.94 (2xpy-CH, 2H, d, J = 2 Hz), 7.04 (2xpy-CH, 2H, d, J = 2 Hz), 7.14 (2xpy-CH, 2H, d, J = 2Hz), 7.18 (2xpy-CH, 2H, d, J = 2Hz), 7.22 (2xpy-CH, 2H, d, J = 2 Hz), 8.24 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 8.74 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.04 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.93 (5xpy-NHCO, 5H, s), 9.96 (py-NHCO, 1H s); MS (FAB), 989 (M-2xCl-H, 0.34).

### EXAMPLE 14

#### Bis-distamycin (Compound 16)

A solution of hexan-1,6-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added

**SUBSTITUTE SHEET**

to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 78% yield. m.p., 210°C;  $^1\text{H-NMR}$ , 1.28 (4,5-suber- $\text{CH}_2$ , 4H, m), 1.57 (3,6-suber- $\text{CH}_2$ , 4H, m), 2.23 (2m7-suber- $\text{CH}_2$ , 4H, tr,  $J = 7$  Hz), 2.63 (2x $\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz), 3.49 (2xCONH $\text{CH}_2$ , 4H, m), 3.80 (2xNCH $_3$ , 6H, s), 3.81 (2xNCH $_3$ , 6H, s), 3.83 (2xNCH $_3$ , 6H, s), 6.88 (2xpy-CH, 2H, d,  $J = 2$  Hz), 6.94 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.05 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.15 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.18 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.23 (2xpy-CH, 2H, d,  $J = 2$  Hz), 8.25 (2xCONH $\text{CH}_2$ , 2H, m), 8.72 [2xC(NH $_2$ ) $_2$ Cl, 4H, s], 9.03 [2xC(NH $_2$ ) $_2$ Cl, 4H, s], 9.86 (2xpy-NHCO, 2H, s), 9.92 (4xpy-NHCO, 4H, s); MS (FAB), 1045 (M-2xCl-H, 0.38).

#### EXAMPLE 15

##### Bis-distamycin (Compound 17)

A solution of octan-1,8-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min. a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 65% yield. m.p., 198-202°C;  $^1\text{H-NMR}$ , 1.26 [(4,5,6,7-seba- $\text{CH}_2$ , 8H, m), 4H, tr,  $J = 6$  Hz], 1.55 [(3,8-seba- $\text{CH}_2$ ), 4H, m], 2.22 (2,9-seba- $\text{CH}_2$ ), 4H, tr,  $J =$

**SUBSTITUTE SHEET**

8 Hz], 2.61 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, tr, J = 6 Hz], 3.48 (2xCONHCH<sub>2</sub>, 4H, m), 3.80 (2xNCH<sub>3</sub>, 6H, s), 3.81 (2xNCH<sub>3</sub>, 6H, s), 3.83 (2xNCH<sub>3</sub>, 6H, s), 6.89 (2xpy-CH, 2H, d, J = 2 Hz), 6.95 (2xpy-CH, 2H, d, J = 2 Hz), 7.05 (2xpy-CH, 2H, d, J = 2 Hz), 7.15 (2xpy-CH, 2H, d, J = 2 Hz), 7.18 (2xpy-CH, 2H, d, J = 2 Hz), 7.22 (2xpy-CH, 2H, d, J = 2 Hz), 8.23 (2xCONHpy-2H, m), 8.65 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.99 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.82 (2xpy-NHCO, 2H, s), 9.91 (4xpy-NHCO, 4H, s); MS (FAB), 1074 (m-2xCl-H, 0.08).

# 10 EXAMPLE 16

## **Bis-distamycin (Compound 18)**

A solution of docosane-1,22-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16 µL, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16 µL, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 73% yield. m.p., 215°C; <sup>1</sup>H-NMR, 1.23 (4,5,...20,21-tetraco-CH<sub>2</sub>, 36H, s), 1.55 (3,22-tetraco-CH<sub>2</sub>, 4H, m), 2.21 (2,23-tetraco-CH<sub>2</sub>, 4H, tr, J = 7 Hz), 2.62 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz], 3.50 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 3.80 (2xNCH<sub>3</sub>, 6H, s), 3.82 (2xNCH<sub>3</sub>, 6H, s), 3.84 (2xNCH<sub>3</sub>, 6H, s), 6.89 (2xpy-CH, 2H, d, J = 2 Hz), 6.94 (2xpy-CH, 2H, d, J = 2 Hz), 7.05 (2xpy-CH, 2H, d, J = 2 Hz), 7.15 (2xpy-CH, 2H, d, J = 2 Hz), 7.19 (2xpy-CH, 2H, d, J = 2 Hz), 7.23 (2xpy-CH, 2H, d, J = 2 Hz), 8.25 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 8.72 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.02 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.83 (2xpy-NHCO, 2H, s), 9.92 (4xpy-NHCO, 4H, s); MS (FAB), 1270 (M-2xCl-H, 0.01).

**SUBSTITUTE SHEET**

EXAMPLE 17**Bis-distamycin (Comp und 29)**

A solution of benzene-1,4-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 77% yield. m.p., >300°C;  $^1\text{H-NMR}$ , 2.63 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz], 3.50 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 6$  Hz), 3.82 ( $2\times\text{NCH}_3$ , 6H, s), 3.86 ( $2\times\text{NCH}_3$ , 6H, s), 3.90 ( $2\times\text{NCH}_3$ , 6H, s), 6.97 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.09 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.15 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.20 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.26 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.38 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 8.10 (aromatic-CH, 4H, s), 8.25 ( $2\times\text{CONHCH}_2$ , 2H, tr,  $J = 6$  Hz), 8.65 [ $2\times\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, s], 9.01 [ $2\times\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, s], 9.95 ( $2\times\text{py-NHCO}$ , 2H, s), 10.03 ( $2\times\text{py-NHCO}$ , 2H, s), 10.57 ( $2\times\text{py-NHCO}$ , 2H, s); ( $\text{CD}_3\text{OD}$ ), 2.71 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 7$  Hz], 3.65 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 7$  Hz), 3.87 ( $2\times\text{NCH}_3$ , 6H, s), 3.91 ( $2\times\text{NCH}_3$ , 6H, s), 3.95 ( $2\times\text{NCH}_3$ , 6H, s), 6.90 ( $2\times\text{ph-CH}$ , 2H, d,  $J = 1.8$  Hz), 6.98 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.07 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.16 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.20 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.34 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 8.04 (AROMATIC-CH, 4H, s); MS (FAB), 1037 ( $\text{M}-2\times\text{Cl-H}$ , 0.05).

**SUBSTITUTE SHEET**

EXAMPLE 18**Bis-distamycin (Compound 30)**

A solution of benzene-1,3-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 68% yield. m.p., 240°C;  $^1\text{H-NMR}$ , 2.61 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz], 3.48 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 6$  Hz), 3.80 ( $2\times\text{NCH}_3$ , 6H, s), 3.86 ( $2\times\text{NCH}_3$ , 6H, s), 3.91 ( $2\times\text{NCH}_3$ , 6H, s), 6.97 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.09 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.16 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.20 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.25 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.38 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.6$  Hz), 7.66 (5-aromatic-CH, 1H, tr,  $J = 7.5$  Hz), 8.10 (4,6-aromatic-CH, 2H, d,  $J^1 = 8$  Hz); 8.21 (2-aromatic-CH, 1H, br, s); 8.21 ( $2\times\text{CONHCH}_2$ , 2H, br, s), 8.58 [ $2\times\text{CY}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 7$  Hz], 3.64 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 7$  Hz), 3.88 ( $2\times\text{NCH}_3$ , 6H, s), 3.90 ( $2\times\text{NCH}_3$ , 6H, s), 3.94 ( $2\times\text{NCH}_3$ , 6H, s), 6.89 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 6.97 ( $2\times\text{ph-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.07 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.20 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.33 ( $2\times\text{py-CH}$ , 2H, d,  $J = 1.8$  Hz), 7.65 (5-aromatic-CH, 1H, tr,  $J = 7.5$  Hz), 8.08 (4,6-aromatic-CH, 2H, d, d,  $J_1 = 7.5$  Hz,  $J_2 = 2$  Hz), 8.47 (2-aromatic-CH, 1H, br, tr,  $J = 2$  Hz); MS (FAB), 1037 ( $-2\times\text{Cl-H}$ , 0.43).

**SUBSTITUTE SHEET**

EXAMPLE 19**Bis-distamycin (Compound 31)**

A solution of benzene-1,2-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 83% yield. m.p., 245°C;  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ ), 2.71 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz], 3.63 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 6$  Hz), 3.87 ( $2\times\text{NCH}_3$ , 6H, s), 3.88 ( $2\times\text{NCH}_3$ , 6H, s), 3.90 ( $2\times\text{NCH}_3$ , 6H, s), 6.89 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 6.91 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 6.97 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.15 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.18 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.24 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.60 ( $2\times\text{m-aromatic-CH}$ , 2H, q,  $J = 3$  Hz), 7.68 ( $2\times\text{o-aromatic-CH}$ , 2H, q,  $J = 3$  Hz); MS (FAB), 1037 ( $\text{m-2xCl-H}$ , 0.65).

EXAMPLE 20**Bis-distamycin (Compound 32)**

A solution of 3,5-pyridine dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from

**SUBSTITUTE SHEET**

methanol and ether. The final product was obtained as a light yellow solid m.p. 250°C in 88% yield. m.p., 250°C; <sup>1</sup>H-NMR, 2.52 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, m], 3.48 (2xCONHCH<sub>2</sub>, 4H, m), 3.81 (2xNCH<sub>3</sub>, 6H, s), 3.85 (2xNCH<sub>3</sub>, 6H, s), 3.88 (2xNCH<sub>3</sub>, 6H, s), 3.90 (2xNCH<sub>3</sub>, 6H, s), 6.96 (2xpy-CH, 2H, m), 7.09 (2xpy-CH, 2H, d, J = 2 Hz), 7.17 (py-CH, 1H, d, J = 2 Hz), 7.19 (2xpy-CH, 2H, d, J = 2 Hz), 7.25 (2xpy-CH, 2H, d, J = 2 Hz), 7.29 (py-CH, 1H, m), 7.40 (py-CH, 1H, m), 7.42 (py-CH, 1H, m), 8.23 (2xCONHCH<sub>2</sub>, 2H, m), 8.25 (3-py-CH, 1H, d, J = 8 Hz), 8.54 (4-py-CH, 1H, m), 8.64 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.99 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.20 (96-py-CH, 1H, m), 9.95 (2xpy-NHCO, 2H, s), 10.04 (2xpy-NHCO, 2H, s), 10.94 (py-NHCO, 1H, s), 11.00 (py-NHCO, 1H, s); (CD<sub>3</sub>OD), 2.72 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz], 3.65 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 3.87 (2xNCH<sub>3</sub>, 6H, s), 3.91 (2xNCH<sub>3</sub>, 6H, s), 3.94 (NCH<sub>3</sub>, 3H, s), 3.954 (NCH<sub>3</sub>, 3H, s), 6.90 (2xpy-CH, 2H, d, J = 2 Hz), 6.98 (2xpy-CH, 2H, d, J = 2 Hz), 7.07 (py-CH, 1H, d, J = 2 Hz), 7.10 (py-CH, 1H, d, J = 2 Hz), 7.15 (2xpy-CH, 2H, d, J = 2 Hz), 7.20 (2xpy-CH, 2H, d, J = 2 Hz), 7.34 (py-CH, 1H, d, J = 2 Hz), 7.41 (py-CH, 1H, d, J = 2 Hz), 8.27 (3-py-CH, 1H, d, J = 8 Hz), 8.44 (4-py-CH, 1H, m), 9.17 (6-py-CH, 1H, m); MS (FAB), 1038 (M-2xCl-H, 0.03).

#### EXAMPLE 21

##### 25 Bis-distamycin (Compound 33)

A solution of pyridine-3,6-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16 μL, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16 μL, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a

**SUBSTITUTE SHEET**



light yellow solid in 74% yield. m.p., 260°C; <sup>1</sup>H-NMR, 2.62 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6Hz), 3.50 (2xCONHCH<sub>2</sub>, 4H, q, J = 6Hz), 3.81 (2xNCH<sub>3</sub>, 6H, s), 3.85 (2xNCH<sub>3</sub>, 6H, s), 3.90 (2xNCH<sub>3</sub>, 6H, s), 6.96 (2xpy-CH, 2H, d, J = 2 Hz), 7.08 (2xpy-CH, 2H, d, J = 2 Hz), 7.16 (2xpy-CH, 2H, d, J = 2 Hz), 7.18 (2xpy-CH, 2H, d, J = 2 Hz), 7.26 (2xpy-CH, 2H, d, J = 2 Hz), 7.39 (2xpy-CH, 2H, d, J = 2 Hz), 8.23 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 8.59 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.87 (4-py-CH, 1H, br, s), 8.98 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.24 (2,6-py-CH, 2H, d, J = 2 Hz), 9.94 (2xpy-NHCO, 2H, s), 10.05 (2xpy-NHCO, 2H, s), 10.83 (2xpy-NHCO, 2H, s); (CD<sub>3</sub>OD), 2.71 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz], 3.64 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 3.87 (2xNCH<sub>3</sub>, 6H, s), 3.99 (2xNCH<sub>3</sub>, 6H, s), 4.02 (2xNCH<sub>3</sub>, 6H, s), 6.88 (2xpy-CH, 2H, s), 6.96 (2xpy-CH, s), 7.07 (2xpy-CH, 2H, s), 7.15 (2xpy-CH, 2H, s), 7.19 (2xpy-CH, 2H, s), 7.35 (2xpy-CH, 2H, s), 8.82 (4-py-CH, 1H, s), 9.17 (2,5-py-CH, 2H, s); MS (FAB), (M-2xCL-H, 0.15).

**EXAMPLE 22****20 Bis-distamycin (Compound 34)**

A solution of pyridine-2,6-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16 μL, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16 μL, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 54% yield. m.p., >260°C; <sup>1</sup>H-NMR, 2.62 [2xCH<sub>2</sub>(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz], 3.50 (2xCONHCH<sub>2</sub>, 4H, m), 3.82 (2xNCH<sub>3</sub>, 6H, s), 3.86 (2xNCH<sub>3</sub>, 6H, s), 3.90 (2xNCH<sub>3</sub>, 6H, s), 6.97 (2xpy-CH, 2H, d, J = 2 Hz), 7.08 (2xpy-CH, 2H, d, J = 2 Hz), 7.15 (2xpy-CH, 2H, d, J = 2

**SUBSTITUTE SHEET**

Hz), 7.18 (2xpy-CH, 2H, d, J = 2 Hz), 7.25 (2xpy-CH, 2H, d, J = 2 Hz), 7.39 (2xpy-CH, 2H, d, J = 2 Hz), 8.23 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 8.56 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.85 (4-py-CH, 1H, tr, J = 2 Hz), 8.96 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.24 (3,5-py-CH, 2H, d, J = 2 Hz), 9.94 (2xpy-NHCO, 2H, s), 10.04 (2xpy-NHCO, 2H, s), 10.81 (2xpy-NHCO, 2H, s); MS (FAB), 1038 (M-2xCL-H, 0.25).

**EXAMPLE 23****Bis-distamycin (Compound 35)**

A solution of trans-1,2-cyclobutane-dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 78% yield. m.p., >230°C; <sup>1</sup>H-NMR, 2.05 (3,4-cyclobutane-CH<sub>2</sub>, 4H, m), 2.60 (2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz), 3.38 (1,2-cyclobutane-CH, 2H, m), 3.49 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 3.79 (2xNCH<sub>3</sub>, 6H, s), 3.84 (2xNCH<sub>3</sub>, 6H, s), 3.85 (2xNCH<sub>3</sub>, 6H, s), 6.88 (2xpy-CH, 2H, d, J = 1.8 Hz), 6.97 (2xpy-CH, 2H, d, J = 1.8 Hz), 7.05 (2xpy-CH, 2H, d, J = 1.8 Hz), 7.17 (2xpy-CH, 2H, d, J = 1.8 Hz), 7.21 (2xpy-CH, 2H, d, J = 1.8 Hz), 7.23 (2xpy-CH, 2H, d, J = 1.8 Hz), 8.22 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 8.55 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.96 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.88 (2xpy-NHCO, 2H, s), 9.94 (4xpy-NHCO, 2H, s); (CD<sub>3</sub>OD), 2.20 (3,4-cyclobutane-CH<sub>2</sub>, 4H, m), 2.71 (2xCH<sub>2</sub>, 4H, tr, J = 7 Hz), 3.49 (1,2-cyclobutane-CH, 2H, m), 3.64 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 7 Hz), 3.87 (2xNCH<sub>3</sub>, 6H, s), 3.89 (2xNCH<sub>3</sub>, 6H, s), 3.90

**SUBSTITUTE SHEET**

(2xNCH<sub>3</sub>, 6H, s), 6.84 (2xpy-CH, 2H, d, J = 2 Hz), 6.89 (2xpy-CH, 2H, d, J = 2 Hz), 6.95 (2xpy-CH, 2H, d, J = 2 Hz), 7.15 (2xpy-CH, 2H, d, J = 2 Hz), 7.18 (2xpy-CH, 2H, d, J = 2 Hz), 7.20 (2xpy-CH, 2H, d, J = 2 Hz); MS (FAB), 1015 (M-2xCl-H, 1.06).

#### EXAMPLE 24

##### Bis-distamycin (Compound 36)

A solution of maleic-dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 33% yield. m.p., >255°C; <sup>1</sup>H-NMR, 2.61 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz), 3.50 (2xCONHCH<sub>2</sub>, 4H, q, J = 6 Hz), 3.82 (2xNCH<sub>3</sub>, 6H, s), 3.85 (2xNCH<sub>3</sub>, 6H, s), 3.87 (2xNCH<sub>3</sub>, 6H, s), 6.97 (2xpy-CH, 2H, tr, J = 2 Hz), 7.07 (2xpy-CH, 2H, d, J = 2 Hz), 7.10 (-CH=CH-, 2H, s), 7.18 (2xpy-CH, 2H, s), 7.24 (2xpy-CH, 2H, d, J = 2 Hz), 7.35 (2xpy-CH, 2H, d, J = 2 Hz), 8.23 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 8.66 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 8.94 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, s], 9.93 (2xpyNHCO, 2H, s), 9.99 (2xpy-NHCO, 2H, s), 10.54 (2xpy-NHCO, 2H, s), (CD<sub>3</sub>OD), 2.72 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, tr, J = 6 Hz), 3.65 (2xCONHCH<sub>2</sub>, 4H, tr, J = 6 Hz), 3.88 (2xNCH<sub>3</sub>, 6H, s), 3.90 (2xNCH<sub>3</sub>, 6H, s), 3.92 (2xNCH<sub>3</sub>, 6H, s), 6.91 (2xpy-CH, 2H, tr, J = 2 Hz), 6.98 (2xpy-CH, 2H, d, J = 2 Hz), 7.09 (-CH=CH-, 2H, s), 7.16 (2xpy-CH, 2H, d, J = 2 Hz), 7.19 (2xpy-CH, 2H, d, J = 2 Hz), 7.33 (2xpy-CH, 2H, d, J = 2 Hz); MS (FAB), 987 (M-2xCl-H, 0.27).

**SUBSTITUTE SHEET**

EXAMPLE 25**Bis-distamycin (Compound 37)**

A solution of fumaroyl-dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent evaporated and the crude product was obtained as a light yellow solid in 67% yield. m.p., >280°C;  $^1\text{H-NMR}$ , 2.61 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz], 3.48 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 6$  Hz); 3.80 ( $2\times\text{NCH}_3$ , 6H, s), 3.84 ( $2\times\text{NCH}_3$ , 6H, s), 3.86 ( $2\times\text{NCH}_3$ , 6H, s), 6.35 ( $-\text{CH}=\text{CH}-$ , 2H, s), 6.84-7.84 ( $12\times\text{py-CH}$ , 12H, m), 8.24 ( $2\times\text{CONHCH}_2$ , 2H, tr,  $J = 6$  Hz), 8.58-9.50 [ $2\times\text{C}(\text{NH}_2)_2\text{Cl}$ , 8H, br, s], 9.93 ( $2\times\text{py-NHCO}$ , 2H, s), 9.97 ( $2\times\text{py-NHCO}$ , 2H, s), 9.98 ( $2\times\text{py-NHCO}$ , 2H, s); ( $\text{CD}_3\text{OD}$ ), 2.66 [ $2\times\text{CH}_2\text{C}(\text{NH}_2)_2\text{Cl}$ , 4H, tr,  $J = 6$  Hz], 3.58 ( $2\times\text{CONHCH}_2$ , 4H, tr,  $J = 6$  Hz), 3.79 ( $2\times\text{NCH}_3$ , 6H, s), 3.82 ( $2\times\text{NCH}_3$ , 6H, s), 3.84 ( $2\times\text{NCH}_3$ , 6H, s), 6.26 ( $-\text{CH}=\text{CH}-$ , 2H, s), 6.83 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 6.87 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 6.91 ( $2\times\text{py-CH}$ , d,  $J = 2$  Hz), 7.13 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.17 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz), 7.27 ( $2\times\text{py-CH}$ , 2H, d,  $J = 2$  Hz); MS (FAB), no M+1 peak.

EXAMPLE 26**Bis-distamycin 35 (Compound 38)**

A solution of trans-5,6-bicyclo[2,2,1]-hept-2-ene dicarbonyl dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of deformyl distamycin (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution

**SUBSTITUTE SHEET**

of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether.

5 The final product was obtained as a light yellow solid in 53% yield. m.p., 260°C;  $^1\text{H-NMR}$ , 1.31 (7-bicyclohept, 1H, s), 1.86 (7-bicyclohept, 1H, d,  $J = 7$  Hz), 2.76 (5-endo-bicyclohept, 1H, d,  $J = 8$  Hz), 2.93 (4-bicyclohept, 1H, s), 3.35 (1-bicyclohept, 1H, s), 3.50 (6-exo-cyclohept, 1H, s), 3.50 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, m], 3.81 (3xNCH<sub>3</sub>, 9H, s), 3.85 (3xNCH<sub>3</sub>, 9H, s), 5.98 (3-bicyclohept, 1H, d, d,  $J = 2.5$  Hz), 6.30 (2-bicyclohept, 1H, d, d,  $J = 2.5$  Hz), 6.86 (py-CH, 1H, d,  $J = 2$  Hz), 6.91 (py-CH, 1H, d,  $J = 2$  Hz), 6.97 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.06 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.13 (py-CH, 1H, d,  $J = 2$  Hz), 7.18 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.19 (py-CH, 1H, d,  $J = 2$  Hz), 7.23 (2xpy-CH, 2H, tr,  $J = 2$  Hz), 8.24 (2xCONHCH<sub>2</sub>, 2H, m), 8.57 [2xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, m] 8.97 [23xC(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, m], 9.88 (py-NHCO, 1H, m), 9.92 (4xpy-CH, 4H, m), 10.11 (Ipy-NHCO, 1H, m); (CD<sub>3</sub>OD), 1.47 (7-bicyclohept, 1H, d,  $J = 8$  Hz), 1.94 (7-bicyclohept, 1H, d,  $J = 8$  Hz), 2.71 [2xCH<sub>2</sub>C(NH<sub>2</sub>)<sub>2</sub>Cl, 4H, d,  $J = 6$  Hz], 2.77 (5-endo-bicyclohept, 1H, d,  $J = 4$  Hz), 3.04 (4-bicyclohept, 1H, s), 3.47 (6-exo-bicyclohept, 1H, s), 3.64 (2xCONHCH<sub>2</sub>, 4H, tr,  $J = 6$  Hz), 3.87 (2xNCH<sub>3</sub>, 9H, s), 3.89 (NCH<sub>3</sub>, 3H, s), 3.90 (2xNCH<sub>3</sub>, 6H, s), 6.08 (3-bicyclohept, 1H, d,  $J = 2.5$  Hz), 6.37 (2-bicyclohept, 1H, d,  $J = 2.5$  Hz), 6.82 (py-CH, 1H, d,  $J = 2$  Hz), 6.83 (py-CH, 1H, d,  $J = 2$  Hz), 6.89 (2xpy-CH, 2H, d,  $J = 2$  Hz), 6.95 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.11 (py-CH, 1H, d,  $J = 2$  Hz), 7.14 (2xpy-CH, 2H, d,  $J = 2$  Hz), 7.17 (3xpy-CH, 3H, tr,  $J = 2$  Hz); MS (FAB), 1053 (M-2xCl-H, 0.21).

10

15

20

25

30

SUBSTITUTE SHEET

EXAMPLE 27**Bis-Lexitropsin (Compound 39)**

A solution of maleic-dichloride (9.28 mg, 0.046 mmol) in 5 mL of tetrahydrofuran was added to a solution of 3-[1-methyl-4-(4-amino-1-methylimidazole-2-carboxamido)imidazole-2-carboxamido]propionamidinium hydrochloride (48 mg, 0.09 mmol) and diisopropylethylamine (Hunig's base, 16  $\mu$ L, 0.09 mmol) in 3 mL of dimethylformamide cooled to 0°C. After 10 min, a solution of Hunig's base (16  $\mu$ L, 0.09 mmol) in 3 mL of THF was added to the reaction solution. The resulting mixture was stirred overnight. The solvent was evaporated and the crude product was recrystallized from methanol and ether. The final product was obtained as a light yellow solid in 73% yield. m.p., >250°C;  $^1\text{H-NMR}$ , 1.86 (2xCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, 4H, q, J = 8 Hz), 3.00 [2xCH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, 4H, tr, J = 8 Hz], 3.30 (2xCONHCH<sub>2</sub>, 4H, m), 3.96 (2xNCH<sub>3</sub>, 6H, s), 4.02 (2xNCH<sub>3</sub>, 6H, s), 7.28 (-CH=CH-, 2H, s), 7.54 (2xim-CH, 2H, s), 7.67 (2xim-CH, 2H, s), 8.52 (2xCONHCH<sub>2</sub>, 2H, tr, J = 6 Hz), 9.43 (2xpy-NHCO, 2H, s), 11.01 (2xpy-NHCO, 2H, s); MS (FAB), 777 (M-2xCl-H, 3.11).

Other compounds shown in Table I were similarly prepared and their analytical and physical data are summarized therein.

25 EXAMPLE 28

Drug-DNA binding constants of the compounds of the present invention were estimated. To 2 mL of Tris-EDTA buffer, pH 8, containing 1.3  $\mu$ M ethidium bromide, calf thymus DNA was added to give a final concentration of 1.35  $\mu$ M. The fluorescence was measured after equilibration for a few minutes, using a Turner model 430 spectrofluorometer (Turner Amsco Instruments, Carpinteria, CA) equipped with a 150 W xenon lamp, at an excitation

**SUBSTITUTE SHEET**

wavelength of 525 nm and an emission wavelength of 600 nm. Aliquots of concentrated drug solutions were added and the fluorescence measured. Controls were performed to show that the drugs themselves did not interfere with the fluorescence measurements at the levels employed. From a plot of the decreased fluorescence of the ethidium-DNA complex with increase dose of drug, the concentration of drug needed to reduce the fluorescence by 50% was determined and used to calculate a relative binding constant for the drug, given the binding constant of ethidium to be  $10^7 \text{ M}^{-1}$  under similar conditions.

The results of binding tests are shown in Table II and in Table III.

#### EXAMPLE 29

Compounds of the present invention were tested for anti-Moloney murine leukemia virus (MLV). The method utilized was adapted from Rowe et al (1970) and Lin et al (1987).

The following materials were utilized in the method:

- Retroviruses; rauscher - ATCC 998  
moloney LT(V) - ATCC 190  
Leukosis-sarcoma complex - ATCC 245
- cells; SC-1 - ATCC CRL 1404  
XC - ATCC CCL 165
- minimum essential medium (eagle) with Hanks Bss, supplemented with 10% fetal bovine serum, 100 IU  $\text{ml}^{-1}$  penicillin G, 100  $\text{ugml}^{-1}$  streptomycin, 2.5  $\text{ugml}^{-1}$  amphotericin B and non-essential amino acids (Sigma M2025).

**SUBSTITUTE SHEET**

- Dulbecco's modified eagles medium, supplemented with 10% fetal bovine with 5% fetal bovine serum, 100 10 ml<sup>-1</sup> penicillin G, 100 ugml<sup>-1</sup> streptomycin and 2.5 ugml<sup>-1</sup> amphotericin B.
- 5        - minimum essential medium (eagle) with earles salt supplemented with 5% fetal bovine serum, 100 10 ml<sup>-1</sup> penicillin G, 100 ugml<sup>-1</sup> streptomycin and non-essential aminoacids (Sigma M2025).
- phosphate buffered saline.
- 10       - crystal violet dye.
- 24 well plates.
- compounds dissolved in DMSO (or water) to 2-20 ugml<sup>-1</sup> then further diluted in 5% FBS-MEM.

15       Stock cell cultures were prepared in the 10% FBS-Dulbecco. To prepare 24 well plates for experiments, 0.8 ml of  $3.5 \times 10^4$  SC<sup>1</sup> cells ml<sup>-1</sup> were added to each well one day in advance. This was using the 5% FBS-MEM. 0.1 ml of each compound dilution, in triplicate, was added to a well in the plate. 0.1 ml of 20-40 p.f.u. of moloney virus was added to each well of the plate. Those plates were shaken on a mechanical shaker at 0, 30 and 60 minutes. They were incubated for 5 days at 37°C in a 5% CO<sub>2</sub> incubator. The medium was removed and plates were subjected to ultraviolet light (175 W cm<sup>2</sup> at surface) for three minutes.

20

25

0.8 ml of  $2 \times 10^5$  XC cells ml<sup>-1</sup> were added to each well using the 10% FBS-Hanks mem. The plates were incubated at 37°C, 5% CO<sub>2</sub> for 4 days, but the medium was replaced after 2 days. The medium was removed, the wells were washed with pbs and 0.25 ml of 0.05% crystal violet

30

**SUBSTITUTE SHEET**



was added to each well for 2 hours. The plates were washed, dried and the plaques counted.

MIC<sub>50</sub> values were calculated using the formula-

$$\frac{\% \text{ inhibition greater than } 50\% - 50\%}{\% \text{ inhibition greater than } 50\% - \% \text{ inhibition less than } 50\%}$$

to give the interpolative values between two dilutions.

The results of the test are shown in Tables IV and V and demonstrate comparative anti-MLV activity between compounds of the present invention and AZT and DDC.

#### 10 EXAMPLE 30

Compounds of the present invention were tested for anti-HIV activity by the National Cancer Institute (NIH, Bethesda). The procedure used by the National Cancer Institute is described in Weislow, O.W. et al, J. Natl. Cancer Inst., Vol. 81, pages 577-586 (1989). NCI uses this procedure to test for agents active against Human Immunodeficiency Virus (HIV) and is designated to detect agents acting at any stage of the virus reproductive cycle. The assay basically involves the killing of T4 lymphocytes by HIV. Small amounts of HIV are added to cells, and a complete cycle of virus reproduction is necessary to obtain the required cell killing. Agents that interact with virions, cells, or virus gene-products to interfere with viral activities will protect cells from cytolysis. The system is automated in several features to accommodate large numbers of candidate agents and is generally designed to detect anti-HIV activity. However, compounds that degenerate or are rapidly metabolized in the culture conditions may not show activity in this screen. All tests are compared with at least one positive (e.g., AZT-treated) control done at the same time under identical conditions. The procedure is set forth below:

**SUBSTITUTE SHEET**

1. Candidate agent is dissolved in dimethyl sulfoxide (unless otherwise instructed) then diluted 1:100 in cell culture medium before preparing serial half-log<sub>10</sub> dilutions. T4 lymphocytes (CEM cell line) are added and after a brief interval HIV-1 is added, resulting in a 1:200 final dilution of the compound. Uninfected cells with the compound serve as a toxicity control, and infected and uninfected cells without the compound serve as basic controls.
2. Cultures are incubated at 37° in a 5% carbon dioxide atmosphere for 6 days.
3. The tetrazolium salt, XTT, is added to all wells, and cultures are incubated to allow formazan color development by viable cells.
4. Individual wells are analyzed spectrophotometrically to quantitate formazan production, and in addition are viewed microscopically for detection of viable cells and confirmation of protective activity.
5. Drug-treated virus-infected cells are compared with drug-tested noninfected cells and with other appropriate controls (untreated infected and untreated noninfected cells, drug-containing wells without cells, etc.) on the same plate.
6. Data are reviewed in comparison with other tests done at the same time and a determination about activity is made.

The test results for five of the active compounds are set forth in the Figures 2-6 and the corresponding Tables VI-X below and test results of the compounds of the present invention are compiled in Table XI.

**SUBSTITUTE SHEET**

Table I. Analytical and physical data on linked  
netropsins and their precursors

	Comp.	Yield(5)	m.p. <sup>a</sup>	Formula	Analysis
5	15	85	210°	C <sub>46</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	16	76°	210°	C <sub>50</sub> H <sub>60</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	17	84	198-202	C <sub>52</sub> H <sub>64</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	18	69	215	C <sub>66</sub> H <sub>92</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	19a	99	305-6°	C <sub>38</sub> H <sub>38</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
10	19b	64	262-8°	C <sub>38</sub> H <sub>46</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	20a	95	278-82°	C <sub>38</sub> H <sub>38</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	20b	78	248-50°	C <sub>38</sub> H <sub>46</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	21a	84.7	289-90°	C <sub>34</sub> H <sub>36</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	21b	58	295°	C <sub>34</sub> H <sub>44</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
15	22a	56.5	250-2°	C <sub>34</sub> H <sub>36</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	22b	85	217°	C <sub>34</sub> H <sub>44</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	23a	88.6	312° (dec)	C <sub>35</sub> H <sub>38</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	23b	68.5	210° (softens)	C <sub>35</sub> H <sub>46</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	24a	59	175°	C <sub>35</sub> H <sub>38</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
20	24b	70.6	204° (softens)	C <sub>35</sub> H <sub>46</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	25a	69	172° (softens)	C <sub>36</sub> H <sub>40</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	25b	77	238° (softens)	C <sub>36</sub> H <sub>48</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	26a	70	165-8°	C <sub>37</sub> H <sub>42</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	26b	46	231°	C <sub>37</sub> H <sub>50</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
25	27a	82.6	189°	C <sub>38</sub> H <sub>44</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	27b	61	201° (softens)	C <sub>38</sub> H <sub>52</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	28a	54	175°	C <sub>38</sub> H <sub>44</sub> N <sub>12</sub> O <sub>6</sub>	C, H, N
	28b	23	198°	C <sub>38</sub> H <sub>52</sub> N <sub>14</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl
	29	77	>300	C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
30	30	68	240	C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	31	83	245	C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	32	88	250	C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	33	74	260	C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	34	54	260	C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
35	35	78	230	C <sub>48</sub> H <sub>60</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	36	33	255	C <sub>46</sub> H <sub>56</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	37	67	280	C <sub>46</sub> H <sub>56</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub>	C, H, N, Cl
	38	53	260	C <sub>51</sub> H <sub>62</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>3</sub>	C, H, N, Cl
	39	73	250	C <sub>34</sub> H <sub>50</sub> N <sub>16</sub> O <sub>6</sub> Cl <sub>2</sub>	C, H, N, Cl

a. Uncorrected.

b. All compounds gave satisfactory elemental analyses within 0.4% of the calculated values and exhibited <sup>1</sup>H-NMR, IR and MS data consistent with the structures.

**SUBSTITUTE SHEET**

TABLE II. Relative binding constants for natural and linked oligopeptides  $R_1CO(CH_2)_nCO-R_1$  to calf thymus DNA determined by ethidium displacement assay.<sup>a</sup>

	Compound	$n^b$	DNA Binding Constant ( $M^{-1}$ )
5	1	--	$1.9 \times 10^7$
	2	--	$0.8 \times 10^7$
	3	0	$5.6 \times 10^7$
	4	1	$3.6 \times 10^7$
	5	2	$7.2 \times 10^7$
10	8	5	$1.2 \times 10^7$
	9	6	$2.5 \times 10^7$
	10	7	$0.9 \times 10^7$
15	11	8	$1.7 \times 10^7$
	12	9	$1.9 \times 10^7$
	13	10	$2.2 \times 10^7$

<sup>a</sup> Based on a binding constant of ethidium of  $10^7 M^{-1}$  under similar conditions of temperature, pH and ionic strength. Binding constant values represent the average of repeat measurements.

<sup>b</sup> Number of  $CH_2$  units in the linker in  $R_1-CO(CH_2)_nCO-R_1$ .

TABLE III. Relative binding constants for cis and trans bis-netropsins to poly(dA-dT) determined by the ethidium displacement assay.<sup>a</sup>

	Compound	$K_{app} (M^{-1})$
30	1	$9.4 \times 10^7$
	2	$6.3 \times 10^7$
	19b	$4.3 \times 10^7$
	20b	$4.9 \times 10^7$
	21b	$4.9 \times 10^7$
35	22b	$3.8 \times 10^7$
	23b	$5.3 \times 10^7$
	24b	$4.4 \times 10^7$
	25b	$5.6 \times 10^7$
	26b	$3.1 \times 10^7$
40	27b	$4.0 \times 10^7$

<sup>a</sup> Based on a binding constant of ethidium of  $9.5 \times 10^6 M^{-1}$  under similar conditions of temperature, pH and ionic strength. Binding constants represent the average of repeat measurements.

**SUBSTITUTE SHEET**

TABLE IV.

Compound	Toxicity, ( $\mu\text{g mL}^{-1}$ )	TD <sub>50</sub>	Activity, MIC <sub>50</sub> ( $\mu\text{g mL}^{-1}$ )	T.I., TD <sub>50</sub> / MIC <sub>50</sub>
5	29	>100.00	3.98	>25.13
	30	>100.00	>50.0	2.0
	31	>100.00	79.63	>1.26
	32	>100.00	15.93	>6.28
	33	>100.00	>100.00	--
10	34	>100.00	22.74	>4.40
	35	83.50	>50.0	1.7
	36	100.00	0.16	625.00
	37	84.29	11.21	7.52
	38	>100.00	22.04	>4.54
15	39	>100.00	>100.00	--
	AZT	>100.00	0.0014	>7.14 $\times 10^5$
	DDC	>100.00	0.74	>135.14

20 TABLE V. Inhibition of Moloney murine leukemia (MLV) associated reverse transcriptase activity by linked.

	<u>Compound</u>	<u>n<sup>a</sup></u>	<u>ID<sub>50</sub><sup>b</sup> (uq/mL) (average ± SD)</u>
25	4	1	39.0 ± 13.9
	4	2	25.2 ± 11.4
	8	5	72.5 ± 7.69
	9	6	21.3 ± 6.1
	10	7	34.2 ± 0.9
30	11	8	20.3 ± 9.2
	12	9	10.3 ± 7.5
	13	10	9.1 ± 6.7
	23b	--	7.0 ± 3.6
	24b	--	30.4 ± 19.3
35	25b	--	21.8 ± 9.2
	26b	--	45.9 ± 11.3
	27b	--	29.1 ± 6.0
	28b	--	63.8 ± 41.0
	Aurintricarboxylic acid		1.42 ± 0.26

40 <sup>a</sup> Number of CH<sub>2</sub> groups in linker in R<sub>1</sub>-CO(CH<sub>2</sub>)<sub>n</sub>CO-R<sub>1</sub>.

<sup>b</sup> 50% inhibitory dose, measured after 120 min incubation of the reaction mixtures. [MLV: lot 804-845-8A; (<sup>3</sup>H-methyl)dTTP at 10  $\mu\text{C}$ ; (specific activity: 30 Curies/mmol) per 250  $\mu\text{L}$  of reaction mixture.]

SUBSTITUTE SHEET

TABLE VI

Results of the compound N,N'-di[1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] terephthalamide dihydrochloride.

5

SUMMARY		DOSE	INFECTED RESPONSE	UNINFECTED RESPONSE
Index	Concentration	(Molar)	% of Control	% of Control
IC50 (Molar)	$> 1.79 \times 10^{-5}$	$5.68 \times 10^{-9}$	39.64	88.62
EC50 (Molar)	$2.08 \times 10^{-6}$	$1.79 \times 10^{-8}$	33.00	92.56
TI50 (IC/EC)	$> 8.59 \times 10^0$	$5.68 \times 10^{-8}$	25.70	94.69
		$1.79 \times 10^{-7}$	29.97	92.72
		$5.67 \times 10^{-7}$	28.86	88.21
		$1.79 \times 10^{-6}$	51.73	136.50
		$5.66 \times 10^{-6}$	134.54	164.46
		$1.79 \times 10^{-5}$	167.25	192.17

10

TABLE VII

Results of the compound N,N'-di[1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] isophthalamide dihydrochloride.

15

SUMMARY		DOSE	INFECTED RESPONSE	UNINFECTED RESPONSE
Index	Concentration	(Molar)	% of Control	% of Control
IC50 (Molar)	$2.84 \times 10^{-4}$	$4.28 \times 10^{-7}$	48.47	116.38
EC50 (Molar)	$3.55 \times 10^{-6}$	$1.35 \times 10^{-6}$	29.25	131.95
TI50 (IC/EC)	$8.00 \times 10^{+1}$	$4.27 \times 10^{-6}$	64.76	123.64
		$1.35 \times 10^{-5}$	117.45	117.76
		$4.26 \times 10^{-5}$	120.68	123.27
		$1.34 \times 10^{-4}$	63.57	142.80
		$4.26 \times 10^{-4}$	0.02	-0.29
		$1.34 \times 10^{-3}$	4.26	10.39

**SUBSTITUTE SHEET**

TABLE VIII

Results of the compound N,N'-di[1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-4pyrrolyl] fumaride dihydrochloride.

5

SUMMARY		DOSE	INFECTED RESPONSE	UNINFECTED RESPONSE
Index	Concentration	(Molar)	% of Control	% of Control
IC50 (Molar)	$>3.30 \times 10^{-5}$	$1.05 \times 10^{-8}$	18.00	83.05
EC50 (Molar)	$1.67 \times 10^{-6}$	$3.32 \times 10^{-8}$	22.96	87.56
TI50 (IC/EC)	$>1.97 \times 10^{-1}$	$1.05 \times 10^{-7}$	26.15	90.26
		$3.31 \times 10^{-7}$	28.00	88.21
		$1.04 \times 10^{-6}$	38.90	83.87
		$3.31 \times 10^{-6}$	96.78	151.83
		$1.04 \times 10^{-5}$	150.24	154.79
		$3.30 \times 10^{-5}$	132.20	138.63

10

TABLE IX

Results of the compound N,N'-di[1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-4pyrrolyl] maleamide dihydrochloride.

15

SUMMARY		DOSE	INFECTED RESPONSE	UNINFECTED RESPONSE
Index	Concentration	(Molar)	% of Control	% of Control
IC50 (Molar)	$1.63 \times 10^{-4}$	$3.04 \times 10^{-7}$	28.75	100.16
EC50 (Molar)	$4.04 \times 10^{-6}$	$9.61 \times 10^{-7}$	32.14	100.02
TI50 (IC/EC)	$4.04 \times 10^{-1}$	$3.03 \times 10^{-6}$	44.03	103.19
		$9.59 \times 10^{-6}$	115.19	112.20
		$3.03 \times 10^{-5}$	114.32	112.05
		$9.58 \times 10^{-5}$	84.02	94.18
		$3.02 \times 10^{-4}$	-0.29	-0.86
		$9.57 \times 10^{-4}$	-0.14	5.40

SUBSTITUTE SHEET

TABLE X

Results of the compound N,N'-di[1-methyl-2-[1-methyl-2-carboxamido(3-propionamidine)-4-pyrrole]-4-pyrrolyl]  
trans 1,2-cyclobutaneamide dihydrochloride.

5

SUMMARY		DOSE	INFECTED RESPONSE	UNINFECTED RESPONSE
Index	Concentration	(Molar)	% of Control	% of Control
IC50 (Molar)	$1.67 \times 10^{-4}$	$2.94 \times 10^{-7}$	42.92	90.95
EC50 (Molar)	$1.39 \times 10^{-6}$	$9.29 \times 10^{-7}$	59.07	102.41
TI50 (IC/EC)	$1.20 \times 10^{+2}$	$2.93 \times 10^{-6}$	77.00	116.36
		$9.28 \times 10^{-6}$	80.26	145.18
		$2.93 \times 10^{-5}$	110.05	174.62
		$9.27 \times 10^{-5}$	90.52	101.95
		$9.29 \times 10^{-4}$	0.04	0.54
		$9.25 \times 10^{-4}$	9.72	8.29

10

TABLE XI

The following Table XI shows the results of anti-HIV-1 data on the oligopeptides of the present invention and their anti-HIV-1 activity is designated as inactive, moderate or active. Compounds 19b, 20b, 21b, 25b, 29, 30,  
 15 32, 34 and 36 are designated as active.

**SUBSTITUTE SHEET**



TABLE XI Anti-HIV-1 Activity

	Compound	IC <sub>50</sub> (μM)	EC <sub>50</sub>	TI <sub>50</sub>	Activity*
5	3	83.5	11.9	7.01	Moderate
	5	75.3	12	6.3	Moderate
	8	64.8	5.3	12.1	Moderate
	9	--	--	--	Inactive
	10	51.1	2.1	24.1	Moderate
10	11	--	--	--	Inactive
	12	57	3.9	1.46	Moderate
	13	78	6.6	11.7	Moderate
	15	41	41	1.0	Inactive
	16	>100	--	--	Inactive
15	17	29	14	2	Moderate
	18	>120	--	--	Inactive
	19b	17.9	1.21	14.8	Active
	20b	284	3.55	80	Active
	21b	33	1.37	24.1	Active
20	22b	199	0.35	566	Active
	23b	9.3	3.44	2.7	Moderate
	24b	257	42.5	6.1	Moderate
	25b	68.2	0.42	161	Active
	26b	168	46.3	3.6	Moderate
25	27b	181	5.6	32.4	Moderate
	29	4.7	0.39	12	Active
	30	140	21	6.6	Active
	32	69	1.6	43	Active
	33	69	9.8	7.0	Moderate
30	34	140	13	11	Active
	35	71	16	4.5	Moderate
	36	207	10.4	19.8	Active
	37	35	--	--	Inactive

\* National Cancer Institute Designation

SUBSTITUTE SHEET

EXAMPLE 31

The following test was an evaluation of compounds according to the invention for inhibition of HBV DNA replication in cell culture.

5      DMVI ASSAY CONTROLS: Untreated Cells, 2',3'-ddC

Test compounds were in the form of powdered materials which were dissolved in DMSO at 30 mg/ml. Compounds were tested for antiviral activity at two concentrations, approximately 10-fold apart, starting in  
10      the range of 100 µg/ml (approximately 100 µM). Compounds were tested for toxicity starting at 300 µg/ml (some were tested starting at 100 µg/ml, and at 50 µg/ml and 10 µg/ml for antiviral activities; 2',3-ddC was requested as a positive control and was tested at the sale  
15      concentrations.

Details of the assay methodology can be found in: Korba and Milman, 1991, Antiviral Res., 15:217. The antiviral evaluations were performed on two separate passages of cells. All wells, in all plates, were seeded  
20      at the same density and at the same time.

Due to inherent variations in the levels of both intracellular and extracellular HBV DNA, only depressions greater than 3.5-fold (for HBV virion DNA) or 3.0-fold (for HBV DNA replication intermediates) from the average  
25      levels for these HBV DNA forms in untreated cells are generally considered to be statistically significant ( $P < 0.05$ ) (Korba and Gerin, manuscript submitted for publication). The levels of integrated HBV DNA in each cellular DNA preparation (which remain constant on a per  
30      cell basis in these experiments) were used, to calculate the levels of intracellular HBV DNA forms, thereby eliminating technical variations inherent in the blot hybridization assays.

**SUBSTITUTE SHEET**

Typical values for extracellular HBV virion DNA in untreated cells range from 50 to 150 pg/ml culture medium (average of approximately 76 pg/ml). Intracellular HBV DNA replication intermediates in untreated cells range from 50 to 100 pg/ $\mu$ g cell DNA (average approximately 74 pg/ $\mu$ g cell DNA). In general, depressions in the levels of intracellular HBV DNA due to treatment with antiviral compounds are less pronounced, and occur more slowly, than depressions in the levels of HBV virion DNA (Korba and Milman, 1991, Antiviral Res., 15:217).

For reference, the manner in which the hybridization analyses were performed for these experiments results in an equivalence of approximately 1.0 pg intracellular HBV DNA/ $\mu$ g cellular DNA to 2-3 genomic copies per cell and 1.0 pg of extracellular HBV DNA/ml culture medium to  $3 \times 10^5$  viral particles/ml.

Toxicity analyses were performed in order to assess whether any observed antiviral effects are due to a general effect on cell viability. The method used was uptake of neutral red dye, a standard and widely used assay for cell viability in a variety of virus-host systems, including HSV and HIV. Details of the procedure are provided in the toxicity table legends.

These studies were performed at the Division of Molecular Virology and Immunology (DMVI) of Georgetown University (GU) and were supported by contract between GU and the National Institute of Allergy and Infectious Diseases (NIAID) as part of the Hepatitis Antiviral Program of the NIAID.

#### TOXICITY EVALUATIONS (TABLE T1)

No significant toxicity (greater than 50% depression of the dye uptake levels observed in untreated cells) was observed for any of the test compounds at the concentrations used for the antiviral evaluations.

**SUBSTITUTE SHEET**

5           No significant toxicity (less than 50% of the dye uptake of untreated cells) was observed for any of the test compounds at the highest concentrations used in the toxicity evaluations. The toxicity of 2',3'-ddC was approximately the same as that previously observed in this culture system (Korba and Gerin, manuscript submitted for publication).

#### ANTIVIRAL EVALUATIONS (TABLES AV1 TO AV3) CONTROLS

10           Within normal variations, levels of HBV virion DNA and intracellular HBV replication intermediates (HBV RI) remained constant in the untreated cells over the challenge period. The positive treatment control, 2,3'-dideoxycytosine (2',3'-ddC), induced significant  
15           depressions of HBV DNA replication at the concentration used (Table AV1). At 9  $\mu$ M 2',3'-ddC, a 90% depression of HBV RI (relative to average levels in untreated cells) is typically observed in this assay system (Korba and Milman, 1991).

#### 20       TEST COMPOUNDS

          Compounds 2, 4-9 and 15 (see pages 68-70) exhibited no significant effect on HBV replication at the concentrations tested.

25           Compound 3 was a weak inhibitor of HBV replication. HBV virion DNA and HBV RI were depressed to a lesser degree than observed following treatment with 2',3'-ddC.

30           Compounds 11 and 12 were moderate inhibitors of HBV replication. HBV virion DNA and HBV RI were depressed to a degree comparable to, but generally less than, that observed following treatment with 2',3'-ddC.

**SUBSTITUTE SHEET**

Compounds 14, 17 and 13 were potent inhibitors of HBV replication. HBV virion DNA and HBV RI were depressed to a degree comparable to, or greater than, that observed following treatment with 2',3'-ddC.

5

Table AV1

Effect of test compounds on HBV replication in 2.2.15 cell cultures

	WELL	TREATMENT	INTRACELLULAR HBV DNA* (pg/ $\mu$ g CELL DNA) MONO. RI		HBV VIRION DNA# (pg/ml CULTURE MEDIUM) DAY 0 DAY 6 DAY 9		
10	114AA	UNTREATED CELLS	2.5	78	57	61	59
	114AB	"	2.2	81	49	62	62
	114BA	"	2.1	70	55	87	92
	114BB	"	2.5	67	57	70	72
15	114AC	2',3'-ddC @ 50 $\mu$ M	1.2	2	57	26	0
	114AD	"	1.4	2	51	28	0
	114BC	"	1.8	1	58	30	0
	114BD	"	1.7	1	59	20	0
20	114AG	2',3'-ddC @ 10 $\mu$ M	2.5	9	63	43	7
	114AH	"	2.5	7	66	47	5
	114BG	"	2.0	8	78	54	6
	114BH	"	2.0	5	68	45	8
25	114AQ	Compound 9 @ 50 $\mu$ M	2.7	70	58	64	62
	114AR	"	2.9	86	69	74	65
	114BQ	"	2.8	66	77	50	71
	114BR	"	2.2	90	68	51	80
30	114AS	Compound 9 @ 10 $\mu$ M	2.3	77	68	71	63
	114AT	"	2.6	80	97	85	81
	114BS	"	2.7	60	60	52	74
	114BT	"	2.7	65	80	72	79
	114AU	Compound 7 @ 50 $\mu$ M	2.7	54	96	67	63
	114AV	"	2.5	55	72	75	68
	114BU	"	2.3	64	81	83	77
	114BV	"	2.3	63	44	69	66

**SUBSTITUTE SHEET**

68

114AW	Compound 7 @ 10 $\mu$ M	2.1	81	75	68	56
114AX	"	2.3	87	92	67	57
114BW	"	2.2	68	60	57	87
114BX	"	2.6	65	55	50	58

5        \* Analysis of intracellular HBV DNA was 24 hours following the 9th day of treatment. DNA in each cell DNA preparation were used to calculate the levels of episomal 3.2Kb HBV genomes (MONO.) and HBV DNA replication intermediates (RI).

10       # A "zero" indicates an undetectable level of HBV DNA, sensitivity cutoff was 0.1 pg/ml.

Table AV2

Effect of test compounds on HBV replication in 2.2.15 cell cultures

15	WELL	TREATMENT	INTRACELLULAR HBV DNA*		HBV VIRION DNA#		
			(pg/ $\mu$ g CELL DNA)		(pg/ml CULTURE MEDIUM)		
			MONO.	RI	DAY 0	DAY 6	DAY 9
20	115AA	Compound 2 @ 50 $\mu$ M	1.3	55	74	68	62
	115AB	"	2.3	64	93	62	74
	115BA	"	2.4	67	77	73	80
	115BB	"	2.4	73	61	83	99
25	115AC	Compound 2 @ 10 $\mu$ M	1.9	84	77	62	100
	115AD	"	2.3	98	91	68	86
	115BC	"	2.3	73	62	61	77
	115BD	"	2.2	91	66	74	56
	115AE	Compound 4 @ 50 $\mu$ M	2.5	94	63	86	97
	115AF	"	2.8	76	66	70	65
	115BE	"	2.1	74	78	64	61
	115BF	"	2.4	73	74	88	85
30	115AG	Compound 4 @ 10 $\mu$ M	2.3	85	77	52	73
	115AH	"	2.8	91	61	55	81
	115BG	"	2.3	54	77	96	62
	115BH	"	2.1	62	64	59	50
35	115AI	Compound 5 @ 50 $\mu$ M	1.8	51	58	82	44
	115AJ	"	2.8	53	56	74	59
	115BI	"	2.2	60	66	90	61
	115BJ	"	2.1	70	55	54	78

**SUBSTITUTE SHEET**

	115AK	Compound 5 @ 10 $\mu$ M	2.3	98	56	94	62
	115AL	"	2.5	70	87	83	80
	115BK	"	2.9	88	68	60	90
	115BL	"	2.9	69	50	81	89
5	115AM	Compound 6 @ 50 $\mu$ M	1.5	66	71	62	95
	115AN	"	1.6	60	72	75	75
	115BM	"	2.6	77	72	55	64
	115BN	"	2.3	80	88	57	91
10	115AO	Compound 6 @ 10 $\mu$ M	1.6	68	56	87	80
	115AP	"	2.8	51	71	61	81
	115BO	"	2.2	50	86	64	97
	115BP	"	2.0	57	96	68	69
15	115AQ	Compound 3 @ 50 $\mu$ M	2.9	27	58	67	16
	115AR	"	2.2	23	58	44	13
	115BQ	"	2.2	26	68	78	19
	115BR	"	2.4	25	81	76	10
20	115AS	Compound 3 @ 10 $\mu$ M	2.8	97	57	79	69
	115AT	"	2.2	86	64	83	74
	115BS	"	2.6	81	74	38	94
	115BT	"	2.5	62	60	97	92
25	115AU	Compound 8 @ 50 $\mu$ M	2.2	75	55	50	55
	115AV	"	2.6	55	77	38	54
	115BU	"	2.5	77	58	48	70
	115BV	"	2.8	80	79	44	75
	115AW	Compound 8 @ 10 $\mu$ M	2.9	58	81	78	59
	115AX	"	2.5	65	63	58	58
	115BW	"	2.1	50	62	58	82
	115BX	"	2.2	66	82	80	87

30 \* Analysis of intracellular HBV DNA was 24 hours following the 9th day of treatment. DNA in each cell DNA preparation were used to calculate the levels of episomal 3.2Kb HBV genomes (MONO.) and HBV DNA replication intermediates (RI).

35 # A "zero" indicates an undetectable level of HBV DNA, sensitivity cutoff was 0.1 pg/ml.

**SUBSTITUTE SHEET**

Table AV3

Effect of test compounds on HBV replication in 2.2.15 cell cultures

5	WELL	TREATMENT	INTRACELLULAR HBV DNA* ( $\mu\text{g}/\mu\text{g}$ CELL DNA) MONO. RI		HBV VIRION DNA# ( $\mu\text{g}/\text{ml}$ CULTURE MEDIUM) DAY 0 DAY 6 DAY 9		
10	116AA	Compound 11 @ 50 $\mu\text{M}$	2.4	8	8	38	2
	116AB	"	2.5	7	69	24	2
	116BA	"	2.3	7	89	48	4
	116BB	"	2.4	9	95	45	3
	116AC	Compound 11 @ 10 $\mu\text{M}$	2.0	58	87	77	30
	116AD	"	1.9	85	85	98	22
	116BC	"	2.2	83	79	74	20
	116BD	"	2.3	61	55	82	25
15	116AE	Compound 15 @ 50 $\mu\text{M}$	2.2	64	55	68	54
	116AF	"	2.1	74	54	70	66
	116BE	"	2.4	67	57	52	44
	116BF	"	2.5	83	75	98	65
20	116AG	Compound 15 @ 10 $\mu\text{M}$	2.3	62	98	42	59
	116AH	"	2.3	58	51	62	70
	116BG	"	2.2	84	87	52	94
	116EH	"	2.5	82	78	60	54
25	116AI	Compound 12 @ 50 $\mu\text{M}$	1.8	11	59	21	9
	116AJ	"	1.0	10	50	19	8
	116BI	"	1.2	9	99	31	11
	116BJ	"	1.1	12	52	29	14
30	116AK	Compound 12 @ 10 $\mu\text{M}$	2.0	57	42	97	55
	116AJ	"	2.0	68	48	98	91
	116BK	"	1.9	58	52	79	54
	116BL	"	1.9	79	84	81	81
	116AM	Compound 14 @ 50 $\mu\text{M}$	1.9	6	71	26	1
	116AN	"	2.0	2	89	27	1
	116BM	"	2.3	3	51	36	1
	116BN	"	2.0	8	92	37	1
35	116AO	Compound 14 @ 10 $\mu\text{M}$	2.5	70	50	110	21
	116AP	"	2.2	54	61	120	25
	116BO	"	2.3	80	55	77	2
	116BP	"	2.1	75	67	87	20
40	116AQ	Compound 17 @ 50 $\mu\text{M}$	2.4	4	63	73	4
	116AR	"	2.7	8	56	68	2
	116BQ	"	2.3	7	66	63	3
	116BR	"	2.4	5	60	89	3

SUBSTITUTE SHEET



71

	116AS	Compound 17 @ 10 $\mu$ M	2.6	43	46	130	55
	116AT	"	2.5	51	45	140	84
	116BS	"	2.1	68	63	65	50
	116BT	"	2.2	62	52	73	84
5	116AU	Compound 13 @ 50 $\mu$ M	2.5	5	50	47	7
	116AV	"	1.8	5	48	52	8
	116BU	"	2.0	6	99	78	6
	116BV	"	2.3	7	88	50	5
10	116AW	Compound 13 @ 10 $\mu$ M	2.2	77	59	130	100
	116AX	"	2.1	78	60	110	54
	116BW	"	2.6	90	94	88	47
	116BX	"	2.4	61	67	69	80

\* Analysis of intracellular HBV DNA was 24 hours following the 9th day of treatment. DNA in each cell DNA preparation were used to calculate the levels of episomal 3.2Kb HBV genomes (MONO.) and HBV DNA replication intermediates (RI).

# A "zero" indicates an undetectable level of HBV DNA, sensitivity cutoff was 0.1 pg/ml.

20

Table T1

Toxicity analysis of test compounds in 2.2.15 cell

		NEUTRAL RED DYE UPTAKE AT INDICATED DRUG CONCENTRATION (% OF CONTROL)				
	PLATE	COMPOUND	300 $\mu$ g/ml	100 $\mu$ g/ml	30 $\mu$ g/ml	10 $\mu$ g/ml
25	27	2',3'-ddC	17 $\pm$ 2	89 $\pm$ 3	100 $\pm$ 1	99 $\pm$ 1
	25	Compound 9	99 $\pm$ 1	100 $\pm$ 2	99 $\pm$ 1	102 $\pm$ 3
	25	Compound 7	98 $\pm$ 2	101 $\pm$ 2	102 $\pm$ 1	101 $\pm$ 2
	25	Compound 4	99 $\pm$ 1	100 $\pm$ 1	101 $\pm$ 1	99 $\pm$ 1
30	26	Compound 3	82 $\pm$ 2	99 $\pm$ 2	99 $\pm$ 1	99 $\pm$ 1
	26	Compound 8	84 $\pm$ 1	99 $\pm$ 2	99 $\pm$ 2	100 $\pm$ 1
	26	Compound 11	84 $\pm$ 2	100 $\pm$ 1	101 $\pm$ 2	99 $\pm$ 2
	26	Compound 15	76 $\pm$ 2	101 $\pm$ 1	101 $\pm$ 2	99 $\pm$ 3
	26	Compound 12	82 $\pm$ 4	101 $\pm$ 1	99 $\pm$ 1	101 $\pm$ 1
35	26	Compound 14	85 $\pm$ 2	99 $\pm$ 1	98 $\pm$ 1	99 $\pm$ 2
	27	Compound 17	82 $\pm$ 1	99 $\pm$ 1	100 $\pm$ 1	100 $\pm$ 2
	27	compound 13	79 $\pm$ 1	98 $\pm$ 1	99 $\pm$ 2	99 $\pm$ 1

SUBSTITUTE SHEET



72





NEUTRAL RED DYE UPTAKE AT INDICATED  
DRUG CONCENTRATION (% OF CONTROL)








	PLATE	COMPOUND	300 $\mu$ g/ml	100 $\mu$ g/ml	30 $\mu$ g/ml	10 $\mu$ g/ml
5	25	Compound 2	96 $\pm$ 1	100 $\pm$ 1	101 $\pm$ 2	102 $\pm$ 1
	25	Compound 5	101 $\pm$ 2	100 $\pm$ 1	99 $\pm$ 2	101 $\pm$ 1
	26	Compound 6	85 $\pm$ 4	101 $\pm$ 1	98 $\pm$ 1	100 $\pm$ 2

Toxicity analyses were performed in 96-well flat bottomed tissue culture plates. Cells for the toxicity analyses were cultured and treated with test compounds with the same schedule as used for the antiviral evaluations. Each compound was tested at 4 concentrations, each in triplicate cultures. Uptake of neutral red dye was used to determine the relative level of toxicity. The absorbance of internalized dye at 510 nm ( $A_{510}$ ) was used for the quantitative analysis. Values are presented as a percentage of the average  $A_{510}$  values ( $\pm$  standard deviations) in 9 separate cultures of untreated cells maintained on the same 96-well plate as the test compounds. The percentage of dye uptake in the 9 control cultures on plate 25 was  $100 \pm 1$ , on plate 26,  $100 \pm 2$ , and on plate 27,  $100 \pm 2$ .

SUBSTITUTE SHEET




Solubility in						
	L	MF and MW	Amount (mg)	DMSO	MeOH	H <sub>2</sub> O
1		C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1108	3.6	good	partial	fair
2		C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1108	4.6	good	good	good
3		C <sub>50</sub> H <sub>58</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1108	3.3	good	good	good
4		C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub> 1109	3.0	good	good	good

		Solubility in				Amount (mg)	MF and MW	L	
		DMSO	MeOH	H <sub>2</sub> O					
5		good	good	good	3.7	C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub> 1109			
6		good	fair	good	3.7	C <sub>49</sub> H <sub>57</sub> N <sub>19</sub> O <sub>8</sub> Cl <sub>2</sub> 1109			
7		good	good	good	3.7	C <sub>48</sub> H <sub>60</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1086			
8		good	good	good	4.5	C <sub>51</sub> H <sub>62</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1175			
9		good	good	good	3.8	C <sub>46</sub> H <sub>56</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1058			
10		good	good	good	1.5	C <sub>46</sub> H <sub>56</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1058			
11		good	good	good	4.2	C <sub>72</sub> H <sub>84</sub> N <sub>27</sub> O <sub>12</sub> Cl <sub>3</sub> 1624.5			
12	-(CH <sub>2</sub> ) <sub>2</sub> -	good	good	good	4.1	C <sub>46</sub> H <sub>112</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1175			


SUBSTITUTE SHEET

		Solubility in			
	L	MF and MW	Amount (mg)	DMSO	H <sub>2</sub> O
13	-(CH <sub>2</sub> ) <sub>6</sub> -	C <sub>50</sub> H <sub>120</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1171	4.8	good	good
14	-(CH <sub>2</sub> ) <sub>8</sub> -	C <sub>52</sub> H <sub>124</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1199	4.2	good	good
15	-(CH <sub>2</sub> ) <sub>22</sub> -	C <sub>66</sub> H <sub>152</sub> N <sub>18</sub> O <sub>8</sub> Cl <sub>2</sub> 1395	4.3	good	partial
16		C <sub>34</sub> H <sub>50</sub> N <sub>16</sub> O <sub>6</sub> Cl <sub>2</sub> 849	3.0	fair	good



Notes

Similar to Compound 1 but prepared by different procedure and with different water of crystallization





RESULTS

5           A wide range of activities was found among the compounds which were tested. All were tested at 50 and 10  $\mu\text{m}$ . The most potent compounds (Compounds 11, 12, 14, 17 and 13) had activities just slightly less than that observed with the control (2',3'-ddC). All exhibited very little toxicity in the confluent cells used for both the antiviral screening and toxicity testing.

**SUBSTITUTE SHEET**

## IN THE CLAIMS:

1. A method for treating a patient infected with a retrovirus, comprising administering to the patient an antiretroviral effective amount of a compound of the formula:



wherein:

A is a moiety bearing a positive charge and of a size which does not inhibit binding of said compound to nucleic acid sequences associated with the cellular action of retroviruses;

R<sub>1</sub> is a moiety derived from a residue of carbonic acid or a residue of a dicarboxylic acid selected from the group consisting of:

- (i) a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p-\text{H}_{2p}-\text{CO}$  where p equals 1 to 22;
- (ii) a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where q equals 2 to 22;
- (iii) a residue of an aromatic dicarboxylic acid;
- (iv) a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  where r equals 3 to 7, optionally fused to one or more three to seven membered C rings; and

**SUBSTITUTE SHEET**

(v) a residue of a cycloalkene dicarboxylic acid of the formula  $-\text{CO}-\text{C}_s-\text{H}_{2s-4}-\text{CO}$  where  $s$  equals 3 to 7;

5        Het is a five membered heterocyclic moiety selected from the group consisting of a pyrrole, an imidazole, a triazole, a pyrazole, a thiazole, a thiophene, a furan and an oxazole;

         x is 0 or 1;

         y is 0, 1, 2 or 3;

10        z is 0, 1, 2 or 3;

$\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $\text{R}_5$  are attached to a ring atom other than carbon and are independently selected from the group consisting of  $\text{C}_1-\text{C}_6$  alkyl and  $-\text{CH}_2-\text{O}-\text{R}_6$ , where  $\text{R}_6$  is a  $\text{C}_1-\text{C}_6$  alkyl;

15        and salts thereof.

2. The method of claim 1, wherein A is a moiety selected from the group consisting of an amidine, a guanidine, secondary ammonium salts, tertiary ammonium salts, quaternary ammonium salts, sulfonium salts and  
20        phosphonium salts.

3. The method of claim 1, wherein  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $\text{R}_5$  are each a  $\text{C}_1-\text{C}_6$  alkyl.

4. The method of claim 1, wherein  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $\text{R}_5$  are the same and are a  $\text{C}_1-\text{C}_6$  alkyl group.

25        5. The method of claim 1, wherein  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$  and  $\text{R}_5$  are each a methoxymethyl.

6. The method of claim 1, wherein  $\text{R}_1$  is



**SUBSTITUTE SHEET**



7. The method of claim 1, wherein  $R_1$  is a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p\text{H}_{2p}-\text{CO}-$  where  $p$  equals 1 to 22.

5 8. The method of claim 1, wherein  $R_1$  is a residue of a dicarboxylic acid selected from the group consisting of: a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where  $q$  equals 2; a residue of an aromatic dicarboxylic acid; and a residue of a cycloalkane dicarboxylic acid of the  
10 formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  where  $r$  equals 3 to 6.

9. The method of claim 1, wherein the compound is  $N,N'$ -di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] terephthalamide dihydrochloride.


15 10. The method of claim 1, wherein the compound is  $N,N'$ -di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] isophthalamide dihydrochloride.

20 11. The method of claim 1, wherein the compound is  $N,N'$ -di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] fumaramide dihydrochloride.


25 12. The method of claim 1, wherein the compound is  $N,N'$ -di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] maleamide dihydrochloride.

30 13. The method of claim 1, wherein the compound is  $N,N'$ -di[1-methyl-2-[1-methyl-2-carboximido(3-propionamidine)-4-pyrrole]-4-pyrrolyl] trans 1,2-cyclobutanamide dihydrochloride.


14. The method of claim 1, wherein the compound is:




and R is



5 15. The method of claim 1, wherein the compound is:




and R is




**SUBSTITUTE SHEET**

81


16. The method of claim 1, wherein the compound is:



and R is




17. The method of claim 1, wherein the compound is:




**SUBSTITUTE SHEET**


and R is



18. The method of claim 1, wherein the compound is:



and R is



5

19. The method of claim 1, wherein said retrovirus is Human Immunodeficiency Virus.

**SUBSTITUTE SHEET**

20. The method of claim 1, wherein the antiretroviral effective dose is in a range of 1 to 200 mg/kg body weight per day.

5 21. the method of claim 1, wherein the compound is administered intravenously or orally.

22. A compound exhibiting activity against retroviruses, represented by the formula:



$$R_2$$

$$R_3$$


10  $(\text{Het}-\text{NHCO})_z-\text{Het}-(\text{CONH})_x-A$

$$R_4$$

$$R_5$$

wherein  $R_1$  is a moiety derived from a residue of a dicarboxylic acid selected from the group consisting of:  
 15 a residue of a  $C_6$  aromatic dicarboxylic acid; a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-C_q-\text{H}_{2q-2}-\text{CO}-$  where  $q$  equals 2; a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-C_r-\text{H}_{2r-2}-\text{CO}$  where  $r$  equals 3 to 6 optionally fused to one or more three to seven C membered rings, and  $A$ ,  $R_2$ ,  
 20  $R_3$ ,  $R_4$ ,  $R_5$  and  $x$ ,  $y$  and  $z$  are as defined in claim 1; and salts thereof.

23. The compound of claim 22, wherein  $R_1$  is



**SUBSTITUTE SHEET**

24. The compound of claim 22, wherein  $R_1$  is




25. The compound of claim 22, wherein  $R_1$  is a dicarboxylic acid residue of cyclopropane.

26. The compound of claim 22, wherein  $R_1$  is a  
5 dicarboxylic acid residue of cyclopentane.


27. The compound of claim 22, wherein  $R_1$  is a dicarboxylic acid residue of cyclohexane.

28. The compound of claim 22, wherein  $R_1$  is





10

29. The compound of claim 22, wherein  $R_1$  is



30. A pharmaceutical composition suitable for  
the treatment of retroviral infections, comprising a  
15 compound of the formula:



20

**SUBSTITUTE SHEET**

wherein:

A is a moiety bearing a positive charge and of a size which does not inhibit binding of said compound to nucleic acid sequences associated with the cellular action of retroviruses;

$R_1$  is a moiety derived from a residue of carbonic acid or a residue of a dicarboxylic acid selected from the group consisting of:

- (i) a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p-\text{H}_{2p}-\text{CO}$  where  $p$  equals 1 to 16;
- (ii) a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where  $q$  equals 2 to 16;
- (iii) a residue of an aromatic dicarboxylic acid;
- (iv) a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  where  $r$  equals 3 to 7 optionally fused to one or more three to six C membered rings; and
- (v) a residue of a cycloalkene dicarboxylic acid of the formula  $-\text{CO}-\text{C}_s-\text{H}_{2s-4}-\text{CO}$  where  $s$  equals 3 to 7;

Het is a five membered heterocyclic moiety selected from the group consisting of a pyrrole, an imidazole, a triazole, a pyrazole, a thiazole, a thiophene, a furan and an oxazole;

$x$  is 0 or 1;

$y$  is 0, 1, 2 or 3;


$z$  is 0, 1, 2 or 3;

$R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are attached to a ring atom other than carbon and are independently selected from the group consisting of  $\text{C}_1-\text{C}_6$  alkyl and  $-\text{CH}_2-\text{O}-\text{R}_6$  is a  $\text{C}_1-\text{C}_6$  alkyl;

and salts thereof, in a pharmaceutically acceptable carrier.

**SUBSTITUTE SHEET**

31. A process for the preparation of a compound of the formula:



wherein:

10 A is a moiety bearing a positive charge and of a size which does not inhibit binding of said compound to nucleic acid sequences associated with the cellular action of retroviruses;

15 R<sub>1</sub> is a moiety derived from a residue of carbonic acid or a residue of a dicarboxylic acid selected from the group consisting of:

- (i) a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p-\text{H}_{2p}-\text{CO}$  where p equals 1 to 16;
- 20 (ii) a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where q equals 2 to 16;
- (iii) a residue of an aromatic dicarboxylic acid;
- 25 (iv) a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  where r equals 3 to 7 optionally fused to a three to seven C membered ring; and
- (v) a residue of a cycloalkene dicarboxylic acid of the formula  $-\text{CO}-\text{C}_s-\text{H}_{2s-4}-\text{CO}$  where
- 30 s equals 3 to 7;

Het is a five membered heterocyclic moiety selected from the group consisting of a pyrrole, an imidazole, a triazole, a pyrazole, a thiazole, a thiophene, a furan and an oxazole;

35

**SUBSTITUTE SHEET**




x is 0 or 1;

y is 0, 1, 2 or 3;

z is 0, 1, 2 or 3;

5  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are attached to a ring atom other than carbon and are independently selected from the group consisting of  $C_1$ - $C_6$  alkyl and  $-CH_2-O-R_6$  is a  $C_1$ - $C_6$  alkyl;

and salts thereof, comprising the steps of:  
reacting a compound of the formula (II)



with a dicarboxylic acid or the formula (III)





15 and converting B to A to form said moiety bearing a positive charge,  
wherein;

x, y and  $R_1$  are as defined above;

20 B is the same as A or is a group with a nitrile, halogen or sulfide substituent; and

X is a halogen, imidazolidine or other reactive moiety.

32. A method for treating a patient infected with Hepatitis B, comprising administering to the  
25 patient an antiretroviral effective amount of a compound of the formula:



**SUBSTITUTE SHEET**

wherein:

A is a moiety bearing a positive charge and a size which does not inhibit binding of said compound to nucleic acid sequences associated with the cellular action of Hepatitis B;

$R_1$  is a moiety derived from a residue of carbonic acid or a residue of a dicarboxylic acid selected from the group consisting of:

- (i) a residue of a dicarboxylic acid of the formula  $-\text{CO}-\text{C}_p-\text{H}_{2p}-\text{CO}$  where  $p$  equals 1 to 22;
- (ii) a residue of an unsaturated aliphatic dicarboxylic acid of the formula  $-\text{CO}-\text{C}_q-\text{H}_{2q-2}-\text{CO}-$  where  $q$  equals 2 to 22;
- (iii) a residue of an aromatic dicarboxylic acid;
- (iv) a residue of a cycloalkane dicarboxylic acid of the formula  $-\text{CO}-\text{C}_r-\text{H}_{2r-2}-\text{CO}-$  wherein  $r$  equals 3 to 7, optionally fused to one or more three to seven membered C rings; and
- (v) a residue of a cycloalkene dicarboxylic acid of the formula  $-\text{CO}-\text{C}_s-\text{H}_{2s-4}-\text{CO}-$  where  $s$  equals 3 to 7;

Het is a five membered heterocyclic moiety selected from the group consisting of a pyrrole, an imidazole, a triazole, a pyrazole, a thiazole, a thiophene, a furan and an oxazole;

$x$  is 0 or 1;

$y$  is 0, 1, 2 or 3

$z$  is 0, 1, 2 or 3

$R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are attached to a ring atom other than carbon and are independently selected from the group consisting of  $\text{C}_1-\text{C}_6$  alkyl and  $-\text{CH}_2-\text{O}-\text{R}_6$ , where  $\text{R}_6$  is a  $\text{C}_1-\text{C}_6$  alkyl;

or a salt thereof.

**SUBSTITUTE SHEET**

1/6




Fig.1

SUBSTITUTE SHEET

2 / 6




Fig.2

SUBSTITUTE SHEET

3 / 6




Fig.3

SUBSTITUTE SHEET

4 / 6




Fig.4

SUBSTITUTE SHEET

5 / 6





Fig.5

6 / 6

Fig.6





## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 94/00427

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D403/14 A61K31/40 C07D401/14 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 13838 (SYNPHAR LAB) 20 August 1992 see the whole document ---	1-32
X	US,A,4 912 199 (LOWN) 27 March 1990 cited in the application Examples 1-3 in columns 7 and 8 ---	1-32
X	J.MED.CHEM. vol. 32, no. 10, 1989 pages 2368 - 2375 LOWN 'Novel Linked Antiviral and Antitumor Agents Related to Netropsin and Distamycin' Page 2370, examples 13a-25a --- -/--	1-32



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

1 December 1994

Date of mailing of the international search report

15. 12. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Gettins, M

# INTERNATIONAL SEARCH REPORT

Int. .onal Application No

PCT/CA 94/00427

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>J.ORG.CHEM.  vol. 56, no. 2 , 1991  pages 786 - 797  RAO 'Sequence-Selective DNA Binding by  Linked Bis-N-methylpyrrole Dipeptides.'  Page 787, examples 3-9  -----</p>	1-32

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 94/00427

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9213838	20-08-92	AU-A- 1224392	07-09-92
US-A-4912199	27-03-90	NONE	

**THIS PAGE BLANK (USPTO)**